



Contents lists available at UGC-CARE

# International Journal of Pharmaceutical Sciences and Drug Research

[ISSN: 0975-248X; CODEN (USA): IJPSPP]

Available online at [www.ijpsdronline.com](http://www.ijpsdronline.com)

## Research Article

# Molecular Docking and QSAR studies for Modeling Antifungal Activity of Triazine Analogues against Therapeutic Target NMT of *Candida albicans*

Sapna Jain Dabade<sup>1,3</sup>, Dheeraj Mandloi<sup>2\*</sup>, Amritlal V. Bajaj<sup>1</sup>, Harsha Atre<sup>3</sup><sup>1</sup>School of Chemical Sciences, Devi Ahilya Vishwavidyalaya, Indore, Madhya Pradesh, India<sup>2</sup>Institute of Engineering and Technology, Devi Ahilya Vishwavidyalaya, Indore, Madhya Pradesh, India<sup>3</sup>Department of Applied Science, SAGE University, Indore, Madhya Pradesh, India

## ARTICLE INFO

### Article history:

Received: 16 July, 2020

Revised: 04 February, 2021

Accepted: 15 February, 2021

Published: 30 March, 2021

### Keywords:

Antifungal, Molecular Docking, QSAR, Triazine


### DOI:

10.25004/IJPSDR.2021.130204

## ABSTRACT

The s-triazine derivatives have been attracting the attention of researchers due to a broad range of biological applications. Present research deals with a combination of genetic algorithm-multiple linear regression (GA-MLR) based quantitative structure-activity relationships (QSAR) modeling and molecular docking as relevant to triazine analogs in an attempt to investigate their role as novel NMT inhibitors of *Candida albicans*. A penta-variant model which assure all validation criteria up to considerable echelon ( $R^2 = 0.792$ ,  $Q^2 = 0.679$  and  $\Delta = 0.603$ ) supplemented by multicollinearity diagnosis by VIF and tolerance data analysis, signaling the robustness of the QSAR model. The descriptors RDF040v, Ds, Mor04m, X4v, and MATS2p in the projected QSAR model have quantified the role of atomic properties such as topology; atomic van der Waals volume, mass, and polarizability execute vital part to modify the antifungal activity of compounds under investigation. Further, a molecular docking simulation study revealed that three compounds, in particular, showed a superior binding affinity with a re-rank score of -142.594, -138.972, -137.540 kcal/mol. Consequently, this study may turn out to be helpful towards the development and optimization of existing antifungal activity of compounds under investigation.

## GRAPHICAL ABSTRACT



## INTRODUCTION

The s-triazine derivatives have been attracting the interest of researchers on account of wide biological activity viz. anti-bacterial, anti-cancer, anti-inflammatory,

anti-viral, anti-malarial fungicidal, anthelmintic, and anti-tubercular.<sup>[1-7]</sup>

Life-threatening, invasive fungal infections especially caused by *Candida* species, are the main cause of morbidity

\*Corresponding Author: Dr. Dheeraj Mandloi

Address: Assistant Professor in Chemistry (Senior Scale), Institute of Engineering and Technology, Devi Ahilya University (NAAC A+ grade State Govt University), Indore MP 452001, India

Email: [dmandloi@ietdavy.edu.in](mailto:dmandloi@ietdavy.edu.in), [dheerajmandloi@rediffmail.com](mailto:dheerajmandloi@rediffmail.com); Orcid id <https://orcid.org/0000-0002-6099-5754>

Tel.: +91-9407894470

**Relevant conflicts of interest/financial disclosures:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Sapna J. Dabade *et al.* This is an open access article distributed under the terms of the Creative Commons Attribution- NonCommercial-ShareAlike 4.0 International License which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

mortality and, thus, causing universal health problems and economic burden.<sup>[8-10]</sup> Large arsenals of available antifungal drugs are not effective due to the issues like drug resistance, narrow antifungal spectrum, poor efficacy, and toxicity. In this framework, the scientific community has been functioning hard to search for novel drug candidate(s) and target.<sup>[11-17]</sup>

It has been evidence from biochemical and genetic studies that enzyme myristoyl CoA: protein N-myristoyl transferase (NMT) plays key responsibility to preserve the viability of *Candida albicans*;<sup>[18]</sup> the principal cause of systematic fungal infections in immune-compromised persons. In this context, NMT has been recognized as a promising target for current investigation.<sup>[19-21]</sup>

Computer-aided drug design (CADD); especially QSAR and molecular docking, has been gaining researchers' attraction to combat the problems associated with traditional methods of drug design and discovery.<sup>[22-23]</sup>

A combination of QSAR and molecular docking studies have been pursued linking chemical structure and pharmacological activities (physical, chemical, and biological properties) quantitatively, along with forecasting the preferred binding orientation of a ligand within the active site of the target protein NMT.<sup>[24-30]</sup> Present research investigation deals with the combination of GA- MLR based QSAR modeling and molecular docking as relevant to triazine analogs in an effort to investigate their role as novel NMT inhibitors of *Candida albicans*. Consequently, this study may be useful for the development and optimization of prevailing antifungal activity of compounds under investigation.

## MATERIALS AND METHODS

The activity data minimum inhibitory concentration (MIC) of the antifungal compounds were taken from the published work<sup>[3]</sup> and converted to-log MIC in a micromolar level. The 2D structures of all the analogues under investigation were first sketched in Chemdraw Ultra version 8.0 software<sup>[31]</sup> then transformed into 3D structures using Chem3D Ultra version 8.0.<sup>[32]</sup> Energy minimization of stated structures was carried out through molecular mechanics (MM2) force field followed by geometry optimized *via* semi-empirical Austin model (AM1) method workable through MOPAC module. Initially, A vast pool of 2D and 3D descriptors (autocorrelation, topological, gateway, RDF, MoRSE, geometrical, constitutional, edge adjacency descriptors subsets) were calculated for the optimized compounds using Parameter Client software.<sup>[33,34]</sup> The descriptors possessing constant numerical values along with those showing high correlation with others were eliminated by using the V-WSP algorithm proposed by Davide and Todeschini<sup>[35]</sup> using NanoBRIDGES" software based on cut-off values of variance (0.0001) and correlation coefficient (0.99).<sup>[35-37]</sup> The hybrid GA-MLR approach linked with error-based fitness function has

been used as the feature selection and model generation tool.<sup>[38-39]</sup>

The robustness of generated models was ensured by applying various validation techniques.<sup>[40-43]</sup> A research report by Roy *et al.* has come out to be worth mentioning in the aforementioned context.<sup>[44]</sup> Therefore, the metrics proposed by Roy *et al.* were also evaluated to check the stability of the QSAR models. Additionally, VIF index, tolerance and Y scrambling analysis were performed to make sure non-existence of the multicollinearity issue in the proposed QSAR model.<sup>[45-46]</sup> Furthermore, to gain insight into ligand-protein interaction, QSAR analysis has been amalgamated with docking studied data under investigation. Molegro virtual docker (MVD) software<sup>[47]</sup> with default parameters were used to perform molecular docking simulation. Proceeding to docking simulation, validation of the software protocol has been performed by RMSD value calculation. The 3D Crystal structure of the target; NMT (1IYL) was retrieved in the MVD from RCSB protein data bank (<https://www.rcsb.org/>). The hydrogen bonds interactions, scoring functions, steric data were investigated for a better understanding of NMT- triazine binding interaction.<sup>[47-48]</sup>

## RESULT AND DISCUSSION

In the current research, a combination of QSAR analysis followed by docking simulation has been carried out with a view to work out the important structural attributes of s-triazine analogs needed for antifungal activity against *Candida albicans*. The structures of 35 triazine derivatives are listed in Fig. 1. Initially, approximately




Fig. 1: Structure of thirty five triazine derivatives under investigation

3000 descriptors were calculated using Parameter Client software.<sup>[33]</sup> The descriptors holding constant numeric values, more than 90% zero values and highly mutually correlated were eliminated as described in the aforesaid section "Materials and Methods". Subsequently, various mono, bi, tri, tetra, and penta-metric QSAR models were generated using genetic algorithm as descriptor screening approach followed by MLR analysis MLR from a large pool of descriptors.<sup>[36-37]</sup> Few best QSAR models among various generated models are presented below, and the statistical parameters to check the quality of these QSAR models are compiled in Table 1.

1. pMIC (C. albicans) = -1.032(± 0.868)E1s + 5.282(± 1.473) - 0.246(± 0.1734) EEig03x + 1.066(± 0.365) EEig08d... Equation 1
2. pMIC (C. albicans) = 0.849(± 0.641) - 11.946(± 5.584) HATS7v + 0.396(± 0.105) RDF155m + 14.513(± 2.249) Ds... Equation 2
3. pMIC (C. albicans) = -0.203(± 1.222) + 0.408(± 0.0999) RDF155m + 11.987(± 2.014) Ds - 0.003(± 0.499) J3D + 0.897(± 0.306) GATS8p... Equation 3
4. pMIC (C. albicans) = -0.721(± 0.836) + 0.911(± 0.302) GATS8p + 11.017(± 2.268) Ds + 0.074(± 0.087) RTp + 0.417(± 0.099) RDF155m... Equation 4
5. pMIC (C. albicans) = -1.845(± 0.820) + 0.342(± 0.064) RDF040v + 14.471(± 2.238) Ds - 0.373(± 0.123) Mor04m - 0.831(± 0.261) X4v + 0.993(± 0.424) MATS2p... Equation 5
6. pMIC (C. albicans) = -2.557(± 0.592) + 15.655(± 2.151) Ds + 2.941(± 0.498) RDF155v - 0.353(± 0.220) Mor05v - 0.795(± 0.222) RDF145v + 1.1823(± 0.541) MATS2v... Equation 6

The regression results (Table 1) show that the tri-parametric (Model I) does not meet the statistical parameters' threshold value. Similarly, Models II-IV seem to be sound as per statistical validation parameters ( $R^2$  and  $R^2_{adj} > 0.5$  along with Y-randomization parameter  $c > 0.6$ ) but not considered as good one due to a lower  $Q^2$  value than 0.5. It is worthy of mentioning that a report published in 2009 by Kunal Roy and co-workers [49] has evidently highlighted the task of validation as a vital aspect of the QSAR model building approach. They have advocated the need to integrate two new parameters  $r^2_m$  and  $r^2_p$  for a stricter parameter of QSAR model validation (particularly when a regulatory discussion is concerned) rather than considering just the conventional validation parameters. Accordingly, conventionally used statistic validation parameters have been add-on with various metrics  $r^2_m$  and  $r^2_p$  validation parameters in the present investigation.

Additionally, Models I-IV are not convincing as per the acceptable QSAR model criteria of Roy *et al.* as the Average  $r^2_m(LOO)$  is less than 0.5.

Penta-variant Models V and VI represented in Equations 5 and 6 respectively satisfied all the aforesaid validation criteria along with various  $r^2_m$  metrics and  $r^2_p$  generated by Roy and Mitra<sup>[44]</sup> (Table 1).

Thus, these two models appear to be statically robust subjected to the green signal from multicollinearity criteria pointed out by Bolboac and Lorentz.<sup>[46]</sup> In this context, to check the multicollinearity among the modeled descriptors, the corresponding statistical finding of VIF and tolerance data are listed in Table 2. High VIF values 7.119 (VIF > 5, and low tolerance value

**Table 1:** Statistical parameters to check the quality of these QSAR models

| Model No.       | $R^2$ | $R^2_{adj}$ | SEE   | $Q^2$ | SDEP  | Average | delta | $r^2_p$   |
|-----------------|-------|-------------|-------|-------|-------|---------|-------|-----------|
| Threshold value | >0.6  | >0.6        | Low   | >0.5  | High  | >0.5    | < 0.2 | $c > 0.6$ |
| Model I         | 0.548 | 0.504       | 0.399 | 0.382 | 0.439 | 0.273   | 0.158 | 508       |
| Model II        | 0.648 | 0.614       | 0.352 | 0.459 | 0.411 | 0.394   | 0.068 | 610       |
| Model III       | 0.686 | 0.644       | 0.337 | 0.469 | 0.407 | 0.412   | 0.045 | 638       |
| Model IV        | 0.694 | 0.652       | 0.334 | 0.483 | 0.402 | 0.426   | 0.040 | 626       |
| Model V         | 0.754 | 0.712       | 0.304 | 0.671 | 0.321 | 0.576   | 0.122 | 0.689     |
| Model VI        | 0.792 | 0.756       | 0.280 | 0.679 | 0.317 | 0.603   | 0.052 | 0.708     |

$R^2$ : coefficient of determination,  $R^2_{adj}$ : Square of adjusted regression coefficient: SEE: Standard error of estimate,  $Q^2$ : Leave-one-out cross-validated coefficient of determination, SDEP: standard deviation error of prediction, Average delta and  $c$  are various metrics for validation developed by Roy and Mitra (2012)

**Table 2:** Collinearity statistical output to check multicollinearity among descriptors (as per models V and VI)

| Collinearity statistics (Model V) |           |       | Collinearity statistics (Model VI) |           |       |
|-----------------------------------|-----------|-------|------------------------------------|-----------|-------|
| Descriptor                        | Tolerance | VIF   | Descriptor                         | Tolerance | VIF   |
| RDF040v                           | 0.221     | 4.523 | Ds                                 | 0.579     | 1.727 |
| Ds                                | 0.630     | 1.586 | RDF155v                            | 0.189     | 5.299 |
| Mor04m                            | 0.571     | 1.751 | Mor05v                             | 0.510     | 1.959 |
| X4v                               | 0.237     | 4.225 | RDF145v                            | 0.140     | 7.119 |
| MATS2p                            | 0.453     | 2.206 | MATS2v                             | 0.569     | 1.758 |



**Table 3:** Observed activity, predicted biological activity, and residual error data of triazine derivatives as per the proposed QSAR model.

| Compound | Observed pMIC | Predicted pMIC | Residuals | Standard residuals |
|----------|---------------|----------------|-----------|--------------------|
| 1        | 3.526         | 3.308          | 0.217     | 0.774              |
| 2        | 3.869         | 4.123          | -0.254    | -0.905             |
| 3        | 5.074         | 4.751          | 0.323     | 1.150              |
| 4        | 4.471         | 4.431          | 0.041     | 0.144              |
| 5        | 4.145         | 4.349          | -0.204    | -0.725             |
| 6        | 3.543         | 3.532          | 0.011     | 0.041              |
| 7        | 3.242         | 3.479          | -0.237    | -0.844             |
| 8        | 3.849         | 3.757          | 0.093     | 0.330              |
| 9        | 4.150         | 4.329          | -0.179    | -0.637             |
| 10       | 4.451         | 4.448          | 0.003     | 0.010              |
| 11       | 3.203         | 3.511          | -0.308    | -1.097             |
| 12       | 3.824         | 3.584          | 0.240     | 0.854              |
| 13       | 3.523         | 3.800          | -0.277    | -0.985             |
| 14       | 4.125         | 3.604          | 0.521     | 1.856              |
| 15       | 3.528         | 3.395          | 0.133     | 0.473              |
| 16       | 4.431         | 4.316          | 0.115     | 0.409              |
| 17       | 3.829         | 4.234          | -0.404    | -1.440             |
| 18       | 3.248         | 3.889          | -0.642    | -2.285             |
| 19       | 4.452         | 4.762          | -0.310    | -1.104             |
| 20       | 4.753         | 4.566          | 0.187     | 0.665              |
| 21       | 3.202         | 3.486          | -0.284    | -1.012             |
| 22       | 3.849         | 3.765          | 0.084     | 0.299              |
| 23       | 4.752         | 4.507          | 0.245     | 0.871              |
| 24       | 3.522         | 3.485          | 0.036     | 0.129              |
| 25       | 4.165         | 4.127          | 0.038     | 0.136              |
| 26       | 3.298         | 3.042          | 0.256     | 0.912              |
| 27       | 4.445         | 4.037          | 0.408     | 1.454              |
| 28       | 3.512         | 3.506          | 0.007     | 0.024              |
| 29       | 3.531         | 3.845          | -0.314    | -1.118             |
| 30       | 5.054         | 4.315          | 0.739     | 2.633              |
| 31       | 4.150         | 4.144          | 0.007     | 0.024              |
| 32       | 4.792         | 4.853          | -0.061    | -0.217             |
| 33       | 3.191         | 3.109          | 0.082     | 0.291              |
| 34       | 3.194         | 3.354          | -0.161    | -0.573             |
| 35       | 3.922         | 4.073          | -0.151    | -0.538             |

**Table 4:** Correlation matrix showing mutual correlations among the variables used in model V


| Descriptor       | RDF040v | Ds     | Mor04m | X4v    | MATS2p | C. albicans pMIC |
|------------------|---------|--------|--------|--------|--------|------------------|
| RDF040v          | 1       |        |        |        |        |                  |
| Ds               | 0.422   | 1      |        |        |        |                  |
| Mor04m           | 0.583   | 0.285  | 1      |        |        |                  |
| X4v              | 0.870   | 0.426  | 0.539  | 1      |        |                  |
| MATS2p           | 0.576   | -0.587 | -0.564 | -0.560 | 1      |                  |
| C. albicans pMIC | 0.440   | 0.642  | -0.051 | 0.249  | -0.177 | 1                |

0.140 (Tolerance < 0.3) indicate collinearity amongst modeled descriptors (Ds, RDF155v, Mor05v, RDF145v, and MATS2v) in Model VI, whereas the non-existence of multicollinearity in generated model V as revealed from statistical parameters of Table 2, i.e., VIF has been found to be less than five and tolerance is greater than 0.3 for all the modeled descriptors. It is apparent from a comparison of various statistical parameters that model V is more robust than other models achieved and discussed above in the present QSAR studies.

Further, In order to ascertain the predictive ability of the developed Model V, pMIC values of all the training set compounds under investigation were calculated through equation 5 and have been compared with the experimental values. As shown in Table 3 the experimental pMIC values are in good agreement with predicted pMIC values as indicated by the low value of standard residual. Further, the plot between predicted and observed pMIC values (Fig. 2) and correlation matrix (Table 4) confirms that the proposed QSAR Model V has fantastic prediction ability.

A combined examination of statistic validation parameters,  $r_m^2$  and  $r_p^2$  metrics validation parameters as per Roy and Mitra,<sup>[44,49]</sup> ( $R^2 = 0.792$ ,  $Q^2 = 0.679$  and  $= 0.603$ ).

VIF and tolerance data analysis enable us to conclude a reasonable agreement of the wide range of validation criteria. Thereby signaling the robustness of the proposed penta-variant QSAR Model V. Therefore, penta-variant

**Fig. 2:** Plot of experimental versus predicted pMIC values of triazine derivatives as generated by the QSAR model V

expression (Equation V) which assures all validation criteria up to considerable echelon, was considered as the best QSAR model for further study.

The developed QSAR model V shows the association between experimental activities considered as a dependent variable and five descriptors, namely RDF040v, Ds, Mor04m, X4v and MATS2p as independent variables in pursuance to modulate antifungal activities of triazine analogs under study. Descriptors inculcated in the best-projected QSAR model encoded the important properties of the molecules pertinent to the antifungal activity. Density descriptor weighted by atomic electro-topological charges (Ds) is a Whim descriptor and provides information regarding the electronic and topological state of the atom in a molecule. A low value of Ds is required in order to increase antifungal activity as it is negatively correlated with activity.<sup>[50]</sup> X4V (connectivity descriptor) is a valence connectivity index chi-4. Tetra-valence Connectivity Index of Electronic Density gives a sign of the nature of bond connectivity between atoms. Connectivity is directly linked with the flexibility of molecules. The higher the connectivity, the more rigid the molecule is, leading to low activity.<sup>[51]</sup> Moran autocorrelation lag 1/weighted by atomic polarizability (MATS2P) is a 2D-autocorrelation descriptor. The positive sign of the regression coefficient of this descriptor indicates elevated activity with a higher value of MATS2P descriptor, and the polarizability plays a vital role in enhancing the activity. The Mor04m (3D-MorSE-signal04/weighted by atomic masses) belongs to the 3D-MorSE descriptors family. 3D-MorSE descriptors provide information regarding the association of activity with mass. The negative sign of Mor04m descriptor illustrates that antifungal activity decreases with increasing molecular mass.<sup>[52-53]</sup>


As shown in equation 5 the descriptor Ds was the most significant descriptor, as indicated by the highest regression coefficient of +14.471. The positive regression coefficient point towards the high value of Ds is vital for elevated antifungal activity. Likewise, negative regression coefficients of another descriptor, i.e., Mor04m, and X4v indicate a lowering in activity with increased numerical values of these descriptors. Correspondingly, the elevated value of the descriptors with a positive regression coefficient, i.e., RDF040v and MATS2p favor the antifungal activity.

The descriptor Ds was found to be the most influential descriptor, as indicated by the highest regression

coefficient of +14.471. Accordingly, topology, atomic van der Waals volume, mass, and polarizability execute a significant role in modulating the antifungal activity of compounds under investigation. Therefore, the substituents that impart the above-mentioned changes in physico-chemical properties included in the proposed model should be attached or removed to the molecules to increase the biological activity.<sup>[50-53]</sup>

With a view to search for better ligand-protein interaction, QSAR analysis has been followed by molecular docking studies using MVD tool. Validation of software protocol has been ensuring by calculating root mean square deviation (RMSD) value. An RMSD value of 1.04 of all the atoms between co-crystallized and docked poses of R64A confirms software reproducibility.

Consequently, all the 35 triazine derivatives were docked into the binding pocket of NMT. The docked energies, hydrogen bond values of the three most active compounds from the thirty-five dataset are provided in Table 5. It was observed (Ref. Table 5) that compound number 5, 32, and 35, in particular, showed a superior binding affinity with a re-rank score of -142.594, -138.972, -137.540 kcal/mol, respectively in comparison to the co-crystallized ligand R64 with a re-rank score of -135.887 kcal/mol. Hydrogen bond interactions of the most active triazole analogs with NMT are illustrated in Figs. 3-5 It was concluded from the docking simulation that three triazine derivatives were strongly bonded through electrostatic, hydrophobic, and hydrogen bond




**Fig. 3:** Hydrogen bond interactions (as shown by blue dotted line) of compound 5 with active amino acids Leu355, and Leu451 of NMT [PDB: 1IYL]


**Table 5:** Molecular docking interactions of triazine derivatives inside the binding pocket of NMT generated by MVD<sup>[47]</sup>.

| Compound No. | Rank score kcal/mol | No of Hydrogen bonds | Amino Acid involved in Hydrogen Bonding | Hydrogen bond Interaction (Å) |
|--------------|---------------------|----------------------|-----------------------------------------|-------------------------------|
| 5            | -137.540            | 03                   | Leu355, and Leu451                      | -2.081                        |
| 32           | -138.972            | 04                   | Asn110, Asp112 and Asn392               | -3.746                        |
| 35           | -142.594            | 05                   | Thr211, Cys393, and Tyr107              | -1.225                        |
| R64A         | 135.887             | 03                   | Asn392, Tyr354,                         | -4.641                        |





**Fig. 4:** Hydrogen bond interactions (as shown by blue dotted line) of compound 35 with active amino acids Thr211, Cys393, and Tyr107 of NMT [PDB: 1IYL]



**Fig. 5:** Hydrogen bond interactions (as shown by blue dotted line) of compound 32 with active amino acids Asn110, Asp112 and Asn392 of NMT [PDB: 1IYL].

interactions and stabilized into the active site of target protein NMT.

## CONCLUSION

Present research findings are associated with a combination of the GA-MLR based QSAR analysis and docking studies, which may help understand the structural necessities for finding novel and selective antifungal drug candidate(s). Consequently, this study may turn out to be useful towards the development and optimization of prevailing antifungal activity of compounds under investigation.

The descriptors RDF040v, Ds, Mor04m, X4v, and MATS2p in the projected QSAR model have quantified the task of atomic properties such as topology, atomic van der Waals volume, mass and polarizability execute vital part to altered the antifungal activity of compounds under investigation. VIF and Tolerance data indicate the non-existence of multicollinearity among the modeled descriptor, and the model is not by chance correlation. Furthermore, molecular docking analysis signifies that compound numbers 5, 32, and 35, in particular, showed a superior binding affinity with a re-rank score of -142.594, -138.972, -137.540 kcal/mol. Combining the GA-MLR based QSAR analysis in association with molecular docking simulation is valuable in understanding the structural requirements for designing novel antifungal drug candidates (s). Accordingly, this research finding may become valuable towards the development of potent antifungals of these congeners.

## ACKNOWLEDGEMENTS

We acknowledge the UGC-SAP (F.540/13/DRSI/2016 (SAP-I)) for providing funds to School of Chemical Sciences, Devi Ahilya University, Indore, India.

## REFERENCES

- Modh RP, De Clercq E, Pannecouque C, Chikhahia KH. Design, synthesis, antimicrobial activity and anti-HIV activity evaluation of novel hybrid quinazoline-triazine derivatives. *Journal of Enzyme Inhibition and Medicinal Chemistry*. 2013; 29 (1): 100-108.
- Modh RP, Patel AC, Chikhahia KH. Design, synthesis, anti-bacterial, and antifungal studies of novel 3-substituted coumarinyl- triazine derivatives. *Heterocyclic. Communication*. 2013; 19(5): 343-349.
- Mewada NS, Shah, DR, Lakum, HP, Chikhahia KH Synthesis and biological evaluation of novel s-triazine based aryl/heteroaryl entities: Design, rationale and comparative study *Journal of the Association of Arab Universities for Basic and Applied Sciences*. 2016; 20: 8-18.
- Singh, UP, Bhat, HR, Gahtori, P. Antifungal activity, SAR and physicochemical correlation of some thiazole-1,3,5-triazine derivatives. *Journal of Medical Mycology* 2012; 22: 134-141.
- Patel RV, Kumari P, Rajani DP, Chikhahia KH, Synthesis and studies of novel 2-(4-cyano-3-trifluoromethylphenylamino)-4-(quinoline-4-yloxy)-6(piperazinyl/piperidinyl )-s-triazines as potential antimicrobial, antimycobacterial and anti-cancer agents. *European Journal of Medicinal Chemistry* 2011; 46: 4354-4365.
- Singh UP, Bhat, HR, Gahtori P. Antifungal activity, SAR and physicochemical correlation of some thiazole-1,3,5-triazine derivatives. *Journal of Medical Mycology* 2010; 22: 134-141.
- Sunduru N, Gupta L, Chaturvedi V, Dwivedi R, Sinha, S, Chauhan PM. Discovery of new 1,3,5-triazine scaffolds with potent activity against *Mycobacterium tuberculosis* H37Rv. *European Journal of Medicinal Chemistry*. 2010; 45: 3335-3345.
- Molero G, Orejas RD, Garcia FN, Monteoliva L, Pla J, Gil C, Perez MS, Nombela C. *Candida albicans*: genetics, dimorphism and pathogenicity. *International Microbiology*. 1998; 1: 95-106.
- Garbino J, Kolarova L, Rohner P, Lew D, Pichna P, Pittet D. Secular trends of candidemia over 12 years in adult patients at a tertiary care hospital. *Medicine*. 2002; 81(6): 425-433
- Hobson R.P. The global epidemiology of invasive *Candida* infections—is the tide turning? *Journal of Hospital Infection*, 2003; 55: 159-168.
- Brown GD, Denning DW, Gow NA, Levitz SM, Netea MG, White TC. Hidden Killers: Human Fungal Infections. *Science Translational Medicine*. 2012; 4(165):1-10.
- Chang YL, Shang JY, Heitman J, Wellington M, Chen YL. New facets of antifungal therapy. *Virulence*. 2017; 8(2): 222-236.
- Mambro, TD, Guerriero I, Aurisicchio I, Magnani M, Marra E. The Yin and Yang of Current Antifungal Therapeutic Strategies: How Can We Harness Our Natural Defenses? *Frontiers in Pharmacology*. 2019;10:1-11
- Fisher MC, Hawkins NJ, Sanglard D, Gurr SJ. Worldwide emergence of resistance to antifungal drugs challenges human health and food security *Science*. 2018; 360(6390): 739-742.
- Daele RV, Spriet I, Wauters J, Maertens J, Mercier T, Hecke SV, Brüggemann R. Antifungal drugs: What brings the future?. *Medical Mycology*. 2019; 57: 28-43.
- Ngo HX, Garneau TS, Green KD. A complex game of hide and seek: the search for new antifungals. *Medicinal Chemistry Communications*. 2016; 7: 1285-1306.
- Mazu TK, Bricker BA, Flores-Rozas H, Ablordeppey SY. The Mechanistic Targets of Antifungal Agents: An Overview. *Mini Reviews in Medicinal Chemistry*. 2016; 16(7): 555-578.
- Weinberg RA, McWherter CA, Freeman K, Wood DC, Gordon JI, Lee SC. Genetic studies reveal that myristoylCoA: Protein N-myristoyltransferase is an essential enzyme in *Candida albicans*. *Molecular Microbiology*. 1995; 16: 241-250.

19. Devadas B, Freeman SK, Mark EZ, Zupec ME, Lu, H.F.; Nagarajan, SR, Kishore NS, Lodge JK, Kuneman DW, McWherter CA, Vinjamoori DV, Getman DP, Gordon JI, Sikorski JA. Design and Synthesis of Novel Imidazole-Substituted Dipeptide Amides as Potent and Selective Inhibitors of *Candida albicans* MyristoylCoA:Protein N-Myristoyltransferase and Identification of Related Tripeptide Inhibitors with Mechanism-Based Antifungal Activity. *Journal of Medicinal Chemistry*. 1997; 40 (16): 2609-2625.
20. Prasad, K.K.; Toraskar, M.P.; Kadam, V.J. N-myristoyltransferase: a novel target. *Mini reviews in Medicinal Chemistry*. 2008; 8(2): 142-149.
21. McCarthy MW, Kontoyiannis DP, Cornely OA, Perfect JR, Walsh TJ. Novel Agents and Drug Targets to Meet the Challenges of Resistant Fungi *Journal of Infectious Diseases*. 2017; 216(3): S474-S483.
22. Wadood, A.; Ahmed, N.; Shah, L.; Ahmad, A.; Hassan, H.; Shams, S. In-silico drug design: An approach which revolutionized the drug discovery process. *Drug. Design Delivery*. 2013; 1(1): 3.1-4.
23. Tang Y, Zhu W, Chen K, Jiang H. New technologies in computer-aided drug design: Toward target identification and new chemical entity discovery. *Drug Discovery Today: Technologies*. 2006; 3(4): 307-313.
24. Cronin MD. Recent Advances in QSAR Studies: Methods and Applications. In: Puzyn T, Leszczynski, Z, Cronin, MD, editors. *Quantitative Structure-Activity Relationships (QSAR)-Application and Methodology Challenges and Advances in Computational Chemistry and Physics*. Heidelberg: Springer; 2010, Vol. 8, p. 3-11.
25. Yee LC, Wei, YC. *Current Modeling Methods Used in QSAR/QSPR (Statistical Modeling of Molecular Descriptors in QSAR/QSPR, First Edition)* Dehmer, M.; Varmuza, K.; Bonchev, D.; Eds., Wiley-VCH Verlag, 2012.
26. Dabade SJ, Mandloi D, Bajaj AV, Dhingra N. GA-SMLR Based QSAR Modeling and Molecular Docking Studies of Bisamidine Derivatives as NMT Inhibitors. *International Journal of Quantitative Structure-Property Relationship*. 2020; 5(4): 1-31.
27. Dabade SJ, Mandloi D, Bajaj AV, Molecular Docking and QSAR Studies of Coumarin Derivatives as NMT Inhibitors: Simple Structural Features as Potential Modulators of Antifungal Activity. *Letters in Drug Design and Discovery*. 2020; 17(10): 1293-1308.
28. Saikia, S.; Bordoloi, M. Molecular Docking: Challenges, Advances and its Use in Drug Discovery Perspective. *Current Drug Targets*. 2019; 20(5): 501-521.
29. Dabade SJ, Mandloi D, Bajaj AV, Thakur A. *In Silico* Evaluation of Inhibitory Potential of Novel Triazole Derivatives Against Therapeutic Target Myristoyl-CoA:Protein N-Myristoyltransferase (NMT) of *Candida albicans*. *Network Modeling Analysis in Health Informatics and Bioinformatics*. 2020; 9(60): 1-17.
30. Silva, D.R., In silico approaches for screening molecular targets in *Candida albicans*: A proteomic insight into drug discovery and development. *European Journal of Pharmacology*. 2019; 842: 64-69.
31. ChemDraw Ultra 8.0.3 Cambridge Soft Chemical Structure Drawing Standard.
32. Chem3D Draw version 8.0.3 Cambridge Soft molecular modeling and analysis tool.
33. Parameter Client, <http://www.vcclab.org/>
34. VCCLAB (Virtual Computational Chemistry Laboratory). <http://www.vcclab.org/> (Accessed January, 2018).
35. Davide, B.; Todeschini, R. A novel variable reduction method adapted from space-filling designs. *Chemometrics and Intelligent Laboratory Systems*. 2014; 136: 147-154.
36. *NanoBRIDGES" software*: Open access tools to perform QSAR and nano-QSAR modeling. *Chemometrics and Intelligent Laboratory Systems* [http://teqip.jdvu.ac.in/QSAR\\_Tools/](http://teqip.jdvu.ac.in/QSAR_Tools/)
37. Aher RB, Ambure P, Gajewicz A, Puzyn T, Roy K. NanoBRIDGES" Software: Open Access Tools to Perform QSAR and Nano-QSAR Modeling. *Chemometrics and Intelligent Laboratory Systems*. 2015; 147: 1-13.
38. Saxena AK, Prathipati P. Comparison of MLR, PLS and GA-MLR in QSAR analysis. *SAR and QSAR in Environmental Research*. 2003; 14: 433-445.
39. Rogers D, Hopfinger A. J. Application of Genetic Function Approximation to Quantitative Structure-Activity Relationships and Quantitative Structure-Property Relationships *Journal of Chemical Information and Modeling*. 1994; 34: 854.
40. Veerasamy R, Rajak H, Jain A, Sivadasan S, Varghese CP, Agrawal RK. Validation of QSAR Models - Strategies and Importance. *International Journal of Drug Design & Discovery*. 2011; 2(3): 511-519.
41. Gramatica, P. Principles of QSAR models validation: internal and external. *QSAR Combinatorial Science*. 2007; 26(5): 694-701.
42. Golbraikh A, Tropsha A. Beware of  $q^2$ ! *J. of Mole. Graph. and Model*. 2002, 20(4), 269-276.
43. Tropsha A, Gramatica P, Gombar VK. The importance of being earnest: validation is the absolute essential for successful application and interpretation of QSPR models. *QSAR Combinatorial Science*. 2003; 22(1): 69-77.
44. Roy K, Mitra I. On the use of the metric  $rm^2$  as an effective tool for validation of QSAR models in computational drug design and predictive toxicology. *Mini-Reviews in Medicinal Chemistry*. 2012; 12(6): 491-504.
45. Chatterjee S, Hadi AS. *Regression Analysis by Examples*. 3rd ed.; Wiley: New York 2000.
46. Bolboac SD, Lorentz J. Quantitative Structure-Activity Relationships: Linear Regression Modelling and Validation Strategies by Example. *Biomath*. 2013; 2: 1-11.
47. Molegro Virtual Docker 6.0. Software.
48. Thomsen R, Christensen MH. MolDock: a new technique for high-accuracy molecular docking. *Journal of Medicinal Chemistry*. 2006; 49(11): 3315-3321.
49. Roy PP, Paul S, Mitra I. Roy, K. On two novel parameters for validation of predictive QSAR models. *Molecules*. 2009; 14: 1660-1701.
50. Todeschini R, Gramatica P. The Whim Theory: New 3D Molecular Descriptors for Qsar in Environmental Modelling. *SAR QSAR Environmental research* 1997; 7(1-4): 89-115.
51. Kier, L.B. & Hall, L.H., *Molecular Connectivity in Structure-Activity Analysis*. RSP-Wiley, Chichester (UK), 1986.
52. Todeschini, R.; Consonni, V. Molecular descriptors for chemoinformatics. In: Mannhold, R., Kubinyi, H., Folkers, G., (Eds.), *Methods and Principles in Medicinal Chemistry*. Weinheim: Wiley - VCH Verlag GmbH & Company, 2009; Vol. 2, p1257.
53. Todeschini R, Consonni V, Mannhold R. Handbook of Molecular descriptors. In: Kubinyi H, Timmerman H. (Eds.) *Methods and Principles in Medicinal Chemistry* Weinheim: Wiley - VCH Germany, 2000; Vol. 11, p 667.

**HOW TO CITE THIS ARTICLE:** Dabade SJ, Mandloi D, Bajaj AV, Atre H. Molecular Docking and QSAR studies for Modeling Antifungal Activity of Triazine Analogues against Therapeutic Target NMT of *Candida albicans*. *Int. J. Pharm. Sci. Drug Res.* 2021;13(2):140-146. DOI: 10.25004/IJPSDR.2021.130204

