



Contribution of NMR Spectroscopy, Mass Spectrometry and X-Ray Diffractometry Techniques to the Characterization of 2-Chloro-8-methyl-3-formylquinoline

M. Faiz Arshad¹, Allam A. Hassan², Abdulmohsen H. Al Rohaimi¹

¹Department of Pharmaceutical Sciences, College of Pharmacy, Shaqra University, Saudi Arabia

²Department of Chemistry, Faculty of Science, Suez University, Egypt

ABSTRACT

Quinoline and its analogues have been known to possess diverse pharmacological activities. In the last three decades 2-chloro-8-methyl-3-formylquinoline has been utilized as building blocks for the development of various quinolines fused heteroaryl rings. In this article we report the complete spectral details of 2-chloro-8-methyl-3-formylquinoline. The techniques involved in the characterization of title compounds are various NMR experiments such as HSQC, COSY and HMBC. Molecular weight was determined by FAB-mass spectrometry. The powder XRD study revealed crystal behavior of the title compound.

Keywords: HSQC, COSY, HMBC, XRD and 2-chloro-8-methyl-3-formylquinoline.

INTRODUCTION

Quinoline ring containing compounds exhibit potent biological activities which have been proven by number of recent reports. [1-3] Among various available intermediates of quinoline, 2-chloro-3-formyl-quinoline **1** and its related aldehyde analogues occupy a prominent space due to its ability to undergo further multiple [a], [b] or/and [c]-annulations leading to formation of wide variety of rings. The applications of these methodologies have yielded a large number of new heteroaryl fused quinoline derivatives such as pyrrolo[3,4-c2

H-pyrano[2,3-*b*]quinoline, tetrazolo[1,5-*a*]quinoline, isothiazolo[5,4-*b*]quinoline, [4-11] and various new approaches for alkaloids synthesis such as camptothecin [12] (Fig. 1). These intermediates have also been exploited immensely for various functional groups interconversions. [13-14] So far the single crystal analysis of 2-chloro-8-methyl-3-formylquinoline is reported by F. Nawaz Khan *et al.* [15], no one yet made any attempt to present spectral details of the compound. As single crystal analysis and spectral applications are complimentary for the determination of structure, we have aimed in this article to present 2D-NMR (HSQC, COSY, HMBC) details, of 2-chloro-8-methyl-3-formylquinoline (Fig. 2) along with powder XRD and softer ionization mass analyses to confirm

the basic structure.




Fig. 1: Synthesis of anticancer natural product Camptothecin starting from 2-chloro-3-formyl-quinoline 1




Fig. 2: Structural formula of 2-chloro-8-methyl-3-formylquinoline and position of H and C designated in HMBC

MATERIAL AND METHODS

All the chemicals required for the synthesis of 2-chloro-8-methyl-3-formylquinoline were procured from E. Merck (Germany). Melting points was determined by the open tube capillary method and is uncorrected. IR spectra were FT-IR spectrophotometer. C, H, N elemental analysis was carried out on Vario EL III CHNOS-Elementar analyzer and was found to be within the range of $\pm 0.4\%$. The title compound 2-chloro-8-methyl-3-formylquinoline needed in the study was prepared as the per the method reported by O. Meth-Cohn *et al.* [16]

*Corresponding author: Dr. M. Faiz Arshad,
Department of Pharmaceutical Sciences, College of Pharmacy, Shaqra University, Al-dawadmi, P.O. BOX 33, Saudi Arabia; Tel.: +00966594082153;
E-mail: drfaizarshad@gmail.com

NMR Analysis

^1H , ^{13}C NMR, COSY, HSQC, and HMBC spectrum of 2-chloro-8-methyl-3-formylquinoline was recorded on a Bruker at 300 MHz Bruker (DPX) spectropin spectrophotometer operating at 300 MHz for ^1H and 75 MHz for ^{13}C equipped with a 5-mm probe, in deuterated DMSO. Tetramethylsilane (TMS) was used as internal standard and the values of chemical shifts δ are expressed in part per million. Splitting patterns are designated as follows: s, singlet; bs, broad singlet; d, doublet; t, triplet.

Mass Analysis

The mass spectra of title compounds was recorded using Fast Atom Bombardment (FAB) technique on JEOL SX 102/DA-6000 Mass Spectrometer/ Data System using Xenon (6kV, 10mA) as the FAB gas. The accelerating voltage was 10kV and the spectra were recorded at room temperature. m-Nitrobenzyl alcohol (NBA) was used as the matrix. The matrix peaks appeared at m/z 136, 137, 154, 289, 307.

X-ray diffractometry (XRD)

In x-ray studies, an automatic x-ray diffractometer (Philips PW 3710; Eindhoven, The Netherlands) equipped with a PW R30 x-ray generator was used. Nickel-filtered Cu α_1 radiation having a wavelength of 1.5106 Å, operating at 35 kW and 20 mA in the range (2θ) of 5 to 70 degrees, was used. X-ray diffractograms were obtained at a scanning rate of 1 degree (2θ) per minute.

Table 1: HSQC details of 2-chloro-8-methyl-3-formylquinoline

^1H (δ) ppm	^{13}C (δ) ppm	Carbon No.
2.69 (s)	17.37	C _{1'}
8.52 (s)	136.91	C ₄
7.69-7.66 (d)	133.30	C ₅
7.46-7.40 (t)	127.26	C ₆
7.63-7.60 (d)	127.56	C ₇
10.45 (s)	188.83	C _{2'}

Table 2: ^1H - ^1H COSY (DQF-COSY) spectrum of 2-chloro-8-methyl-3-formylquinoline

Position of (H)	^1H -NMR (δ)	COSY
1'	2.69	--
4	8.52	--
5	7.69-7.66	H-6
6	7.46-7.40	H-5, H-7
7	7.63-7.60	H-6
2'	10.45	---

RESULTS AND DISCUSSION

HSQC

The sequential assignment of protons and carbon atoms of 2-chloro-8-methyl-3-formylquinoline was made with the help of HSQC experiment (Table 1 and Figure 3). Starting with the easily distinguishable methyl protons $\delta_{\text{H}} = 2.69$ (s) showed correlation with $\delta_{\text{C}} = 17.37$ attributed at position 1'. The next singlet proton $\delta_{\text{H}} = 8.52$ showed correlation with the carbon $\delta_{\text{C}} = 136.91$ attributable at position 4'. Similarly, the proton $\delta_{\text{H}} = 7.69-7.66$ (d, $J = 8.1$ Hz) exhibited correlation with carbon $\delta_{\text{C}} = 133.30$ assignable at position 5. Another doublet was observed at $\delta_{\text{H}} = 7.63-7.60$ (d, $J = 8.1$) which showed correlation with carbon $\delta_{\text{C}} = 127.56$ attributable at position 7. A triplet was observed at chemical shift value $\delta_{\text{H}} = 7.46-7.40$ (t, $J = 7.65$) assignable at position 6 and correlated with the carbon $\delta_{\text{C}} = 127.26$. The highly distinguishable aldehydic protons $\delta_{\text{H}} = 10.45$ (s) showed correlation with carbon at $\delta_{\text{C}} = 188.83$ assignable at position 2'.




Fig. 3: HSQC- Contour plot of 2-chloro-8-methyl-3-formylquinoline




Fig. 4: COSY-DQF-Contour plot of 2-chloro-8-methyl-3-formylquinoline




Fig. 5: HMBC-Contour plot of 2-chloro-8-methyl-3-formylquinoline




Fig. 6: FAB-MS spectrum of 2-chloro-8-methyl-3-formylquinoline




Fig. 7: Powder XRD diffractogram of 2-chloro-8-methyl-3-formylquinoline

Table 3: HMBC correlation data of compound 2-chloro-8-methyl-3-formylquinoline

Protons	Carbon (ppm)	C1'	C2	C3	C4	C5	C6	C7	C8	C9	C10	C2'
H-1'	2.69	--	--	--	--	--	--	3JCH	2JCH	3JCH	--	--
H-4	8.52	--	3JCH	2JCH	--	3JCH	--	--	--	3JCH	2JCH	3JCH
H-5	7.69-7.66	--	--	--	3JCH	--	2JCH	3JCH	--	3JCH	2JCH	--
H-6	7.46-7.40	--	--	--	--	2JCH	--	2JCH	3JCH	--	3JCH	--
H-7	7.63-7.60	3JCH	--	--	--	3JCH	2JCH	--	2JCH	3JCH	--	--
H-2'	10.45	--	3JCH	2JCH	3JCH	--	--	--	--	--	3JCH	--

¹H-¹H COSY (DQF-COSY)

In ¹H-¹H COSY (DQF-COSY) experiment (Figure 4), the proton exhibited chemical shift values $\delta_H = 2.69$ did not correlate with any other proton. At position 4, the proton having chemical shift value $\delta_H = 8.52$ did not show any correlation. The proton positioned at 5, owing chemical shift value $\delta_H = 7.69-7.66$ showed correlation with H-6. Similarly, the number six proton displayed correlation with two different protons positioned at 5 and 6. The H-7 chemical shift value $\delta_H = 7.63-7.60$ exhibited correlation with H-6 only. The 2' did not show any correlation (Table 2).

HMBC

In HMBC two bond correlations (2JCH) are almost always found, the three bond correlation (3JCH) are occasionally absent (Figure 5). As proceeding parallel to the carbon column we got the following results (Table 3).

1. The H-1' protons ($\delta_H = 2.69$) showed correlation with C7 (3JCH), C8 (2JCH) and C9 (3JCH) carbon atom.

2 The H-4 proton ($\delta_H = 8.52$) displayed multiple correlation such as C2 (3JCH), C3 (2JCH), C5 (3JCH), C9 (3JCH), C10 (2JCH) and C2'(3JCH) respectively.

3 Similarly H-5 proton exhibited correlation with carbon atoms C4 (3JCH), C6 (2JCH), C7 (3JCH), C9 (3JCH) and C10 (2JCH).

4 The H-6 proton correlated with carbon atoms C5 (2JCH), C7 (2JCH), C8 (3JCH) and C10 (3JCH) respectively.

5 The multiple correlation was observed between H-7 proton ($\delta_H = 7.63-7.60$) and carbon atoms C1' (3JCH), C5 (3JCH), C6 (2JCH), C8 (2JCH), and C9 (3JCH) respectively.

6 The aldehydic proton also displayed multiple correlation with the carbon atoms C2 (3JCH), C3 (2JCH), C4 (3JCH) and C10 (3JCH) respectively. As proceeding parallel to the proton column we got the similar results e.g. C1' carbon atom showed only one correlation with proton H-7 (3JCH). Similarly C2 exhibited two correlation with protons H-4 (2JCH) and H-2' (3JCH).

FAB-MS

The mass spectrum of 2-chloro-8-methyl-3-formylquinoline using FAB-MS registered M^+ peak at m/z 205 (100 %) and $M+2$ peak at m/z 207 (30 %) due presence of chlorine atom (Figure 6), the spectrum also shows $M+H$ peak at m/z 206 (10 %). A small peak due loss of H^+ ion, was appeared at 204 (20 %) and peak at m/z 176 may be attributed due to loss of -CHO from the molecular ion. A fragment peak at m/z 169 was also observed in the spectrum which expected to appear due to loss of HCl from the molecular ion.

Powder XRD

Figure 7 shows the powder XRD patterns of powder 2-chloro-8-methyl-3-formylquinoline. The XRD diffractogram of 2-chloro-8-methyl-3-formylquinoline contained sharp peaks indicating good crystallinity. The height of peaks varies with different 2θ values. This diffractogram may serve as a finger print for the characterization of this compound.

In the article we have presented correlation spectroscopic details of 2-chloro-8-methyl-3-formylquinoline. The HMBC studies indicate that all H-5, H-6 H-7 proton shows multiple correlation with carbon atoms.

REFERENCES

1. Patel D, Maulik PS, Patel V, Singh R. BCR ABL Kinase Inhibitors for Cancer Therapy. *Internat. J. Pharm. Sci. Drug Res.* 2010; 2(2): 80-90.
2. Bhargu B, Pathak D, Siddiqui N, Alam MS, Ahsan W. Search for Biological Active Isatins: A Short Review. *Internat. J. Pharm. Sci. Drug Res.* 2010; 2(4): 229-235.
3. Desai UH, Patwari AH, Maradiya JK, Sathawara MK, Suhagia BN, Rathod IS. RP-HPLC Method for Simultaneous Estimation of Ciprofloxacin and Dexamethasone in Eye/Ear Drops. *Internat. J. Pharm. Sci. Drug Res.* 2013; 5(2): 62-66.
4. Nyerges M, Pintér A, Virányi A, Blaskó G, Tőke L. Synthesis of pyrrolo[3,4-*c*]quinolines by 1,5-electrocyclisation of non-stabilized azomethine ylides. *Tetrahedron.* 2005; 61: 8199-8205.
5. Cziaky Z, Szabó Z. Synthesis of 2*H*-pyrano[2,3-*b*]quinolines. Part II. Preparation and 1H -NMR investigations of 4-hydroxy-2-methyl-3,4-dihydro-2*H*-pyrano[2,3-*b*]quinolines. *J. Hetero. Chem.* 1995; 32: 755-760.
6. Bawa S, Kumar S. Synthesis of Schiff's bases of 8-methyl-tetrazolo[1,5-*a*]quinoline as potential anti-inflammatory and antimicrobial agents. *Indian J. Chem.* 2009; 48B: 142-145.
7. Eswaran S, Adhikari AV, Kumar RA. New 1, 3-oxazolo[4,5-*c*]quinoline derivatives: Synthesis and evaluation of antibacterial and antituberculosis properties. *Euro. J. Med. Chem.* 2010; 45: 957-966.
8. Srivastava A, Chandra A, Singh RM. Isothiazolo-fused quinoline analogues : Synthesis of isothiazolo[5,4-*b*]quinoline and their oxidation product, 3[2*H*]-one-1,1 dioxidoisothiazolo[5,4-*b*]quinoline from 2-chloro-3-formylquinoline. *Indian J. Chem.* 2007; 46B: 303-307.
9. Amaravathi M, Babu MM, Chandramouli G. Synthesis of meso-tetrakis (2-chloroquinolin-3-yl) porphyrins. *Arkivoc.* 2007; (i): 148-153.
10. El-Baih FEM, Al-Aqeel SI, El-Sayed OA, Al-Hazimi HM. Synthesis and spectroscopic properties of some pyrimido[4,5-*b*]quinoline derivatives. *J. King Saud. Univ.* 2005; 18: 1-17.
11. Fused pyridine and pyrazine derivatives as kinase inhibitors. 2010 WO/2010/092340.
12. Thomas CJ, Rahier NJ, Hecht SM. Camptothecin: Current perspectives. *Bioorg. Med. Chem.* 2004; 12: 1585-1604.
13. Srivastava A, Singh RM. Vilsmeier-Haack reagent: A facile synthesis of 2-chloro-3-formylquinoline from N-arylacetamide and transformation into different functionalities. *Indian J. Chem.* 2005; 44B: 1886-1875.
14. Bouraiou A, Menasra H, Debache A, Rhouati S, Belfaitah A. Synthesis of some functionalized quinolyl aziridine derivatives. *J. Soc. Alger. Chim.* 2006; 16: 171-183.
15. Khan FN, Subashini R, Kushwaha A K, Hathwar VR, Ng SW. 2-Chloro-8-methyl quinoline-3-carbaldehyde. *Acta Cryst.* 2009; E65E: o2722.
16. Meth-Cohn O, Narine B, Tarnowski B. A versatile new synthesis of quinoline and related fused pyridines. Part-5. The synthesis of 2-chloroquinoline-3-carboxaldehyde. *J Chem Soc. Perkin Trans.* 1981; I: 1520-1530.