

International Journal of Pharmaceutical Sciences and Drug Research

2015; 7(5): 417-420



Research Article

ISSN: 0975-248X
CODEN (USA): IJPSP

Synthesis, Characterization and Antimicrobial Evaluation of Some Tetrahydroquinazoline Derivatives of Benzo[b]thiophene

R. Bhatt, S. Sharma, A. K. Patidar, K. K. Rathore, R. C. Senwar, A. Mehta*

Synthetic Organic Chemistry Laboratory, Department of Chemistry, M. L. Sukhadia University,
Udaipur-313001, Rajasthan, India

ABSTRACT

In the present investigation, synthesis of 4-(1-(3-chlorobenzo[b]thiophene-2-carbonyl)-1H-indol-3-yl)-7, 7-dimethyl-3, 4, 7, 8-tetrahydroquinazoline-2, 5(1H, 6H)dione (**3a-d**) is described. These unreported compounds are obtained starting from indole-3-aldehyde and 3-chloro-1-benzothiophene-2-carbonyl chloride (**1a-d**). A series of compounds (**3a-d**) were prepared. The structures of the newly synthesized compounds were elucidated on the basis of IR and ¹H NMR spectral data of the compounds. All the compounds have been screened for antibacterial activity against *S. aureus*, *E. coli*, and *P. aeruginosa* and for antifungal activity against *Candida albicans*, *A. niger*, and *A. clavatus* using broth dilution method and they were found to exhibit good to moderate antimicrobial activity.

Keywords: 3-chlorobenzo[b]thiophene-2-carbonyl, Synthesis, Antimicrobial activity.

INTRODUCTION

Thiophene ring fused with benzene nucleus is known important pharmacophore that occurs frequently in medicinal chemistry literature. A variety of reports regarding synthetic studies of benzo[b]thiophene derivatives have been presented due to their chemical and biological interests. Benzo[b]thiophene compounds are widely used as antibacterial [1-7], antifungal [8-12], and data, benzo[b]thiophene nucleus has attracted the attention of medicinal chemistry due to its well known anticancer activity and many substituted derivatives have recently earned great interests in chemotherapy as

antitumor drugs. Beside the currently established drugs Zileuton.


Benzo[b]thiophene derivatives are associated with diverse biological activities viz. antiviral [13], antitumor [14-16], anti-inflammatory [17-22], antiallergy [23], antimalarial [24], antimicrobial [25-30] and analgesic. [31] Because of these widespread activities, their synthesis was continually the subject of much research and is still pursued due to new emerging methods. A novel series of benzo[b]thiophen-3-ylmethylidene derivatives was comp good antimicrobial activity.

MATERIAL AND METHODS

The reagent grade chemicals were purchased from commercial sources and purified either by distillation or recrystallization before use. Purity of the compounds was checked on silica gel G TLC plates of 2 mm thickness using n-hexane and ethyl acetate as solvent system. The visualization of spot was carried out in an iodine chamber. The IR spectra of the compounds were

*Corresponding author: Dr. Anita Mehta,
Synthetic Organic Chemistry Laboratory, Department
of Chemistry, M. L. Sukhadia University, Udaipur-
313001, Rajasthan, India;
E-mail: anita.mehta83@yahoo.co.in
Received: 02 May, 2015; Accepted: 21 July, 2015

recorded in the 4000-450 cm^{-1} ranges using KBr discs on FTIR IR RX1 Perkin Elmer Spectrophotometers and NMR were recorded on a Bruker DRX-300 MHz spectrometer ($\text{CDCl}_3/\text{DMSO}$) using TMS as an internal standard.



Synthesis of 1-[(3-chloro-1-benzothiophen-2-yl) carbonyl]-1H-indole-3-carbaldehyde (2a)

A mixture of indole-3-aldehyde (0.01mol) and 3-chloro-1-benzothiophene-2-carbonylchloride (1a-d) (0.01mol) in presence of base were dissolved in ethanol and refluxed for 12 hours at room temperature with stirring. Completion of reaction was monitored using TLC and mixture was kept overnight at room temperature. The precipitate was filtered, washed, dried and recrystallized from ethanol.

IR (KBr) cm^{-1} : 3076(C-H str., Aromatic ring), 1682(C=O str., Amide group), 687(C-S-C str., Benzo[b] thiophene ring), 752(C-Cl str.); ^1H NMR (DMSO-d_6) δ : 7.31-8.19(9H, m, Ar-H), 4.85(1H, s, =C-H, indol ring), 9.65(1H, s, CHO).

1-[(3, 6-dichloro-1-benzothiophen-2-yl) carbonyl]-1H-indole-3-carbaldehyde (2b)

IR (KBr) cm^{-1} : 3074(C-H str., Aromatic ring), 1680(C=O str., Amide group), 682(C-S-C str., Benzo[b] thiophene ring), 758(C-Cl str.); ^1H NMR (DMSO-d_6) δ : 7.31-8.21(7H, m, Ar-H), 4.75(1H, s, =C-H, indol ring), 9.74(1H, s, CHO).

1-[(3-chloro-6-fluoro-1-benzothiophen-2-yl) carbonyl]-1H-indole-3-carbaldehyde (2c)

IR (KBr) cm^{-1} : 3079(C-H str., Aromatic ring), 1687(C=O str., Amide group), 689(C-S-C str., Benzo[b] thiophene ring), 759(C-Cl str.), 1153(C-F str.); ^1H NMR (DMSO-d_6) δ : 7.29-8.15(7H, m, Ar-H), 4.75(1H, s, =C-H, indol ring), 9.52(1H, s, CHO).

1-[(3-chloro-6-hydroxy-1-benzothiophen-2-yl) carbonyl]-1H-indole-3-carbaldehyde (2d)

IR (KBr) cm^{-1} : 3081(C-H str., Aromatic ring), 1680(C=O str., Amide group), 681(C-S-C str., Benzo[b] thiophene ring), 753(C-Cl str.), 3384 (OH str.); ^1H NMR (DMSO-d_6) δ : 7.22-8.19(7H, m, Ar-H), 4.78(1H, s, =C-H, indol ring), 9.21(1H, s, CHO), 5.82(1H, s, O-H).

Synthesis of 4-(1-(3-chlorobenzo[b]thiophene-2-carbonyl)-1H-indol-3-yl) 7, 7-dimethyl-3, 4, 7, 8-tetrahydroquinazoline-2, 5(1H, 6H) dione (3a)

A mixture of compound (2a-d) (0.01mol), 5,5-dimethylcyclohexane-1, 3-dione (0.01mol) and urea (0.012mol) were taken in 100 ml RB flask, and stirred at 100-110°C for 3 hr. Then the mixture was cooled to room temperature. The solid obtained was washed with ethanol and recrystallized to afford (3a-d).

IR (KBr) cm^{-1} : 3080(C-H str., Aromatic ring), 2959(C-H str., -CH₃ group), 3258(N-H str.), 681(C-S-C str., Benzo[b] thiophene ring), 742 (C-Cl str.), 1690(C=O str.); ^1H NMR (DMSO-d_6) δ : 6.91-8.10(8H, m, Ar-H), 1.32(6H, s, Ar-CH₃), 8.56(2H, s, -NH quinazoline ring), 5.56(1H, s, =CH), 1.96(2H, t, CH₂, quinazoline ring), 1.32(2H, t, CH₂, quinazoline ring).

4-(1-(3-chlorobenzo[b]thiophene-2-carbonyl)-1H-indol-3-yl) 7, 7-dimethyl-3, 4, 7, 8-tetrahydroquinazoline-2, 5(1H, 6H) dione (3b)

IR (KBr) cm^{-1} : 3056(C-H str., Aromatic ring), 2958(C-H str., -CH₃ group), 3256(N-H str.), 692(C-S-C str., Benzo[b] thiophene ring), 749 (C-Cl str.), 1686(C=O str.); ^1H NMR (DMSO-d_6) δ : 6.93-8.12(7H, m, Ar-H), 1.32(6H, s, Ar-CH₃), 8.61(2H, s, -NH quinazoline ring), 5.59(1H, s, =CH), 1.98(2H, t, CH₂, quinazoline ring), 1.36(2H, t, CH₂, quinazoline ring).

4-(1-(3-chlorobenzo[b]thiophene-2-carbonyl)-1H-indol-3-yl) 7, 7-dimethyl-3, 4, 7, 8-tetrahydroquinazoline-2, 5(1H, 6H) dione (3c)

IR (KBr) cm^{-1} : 3057(C-H str., Aromatic ring), 2952(C-H str., -CH₃ group), 3265(N-H str.), 684(C-S-C str., Benzo[b] thiophene ring), 749 (C-Cl str.), 1686(C=O str.), 1157(C-F str.); ^1H NMR (DMSO-d_6) δ : 6.92-8.09(7H, m, Ar-H), 1.30(6H, s, Ar-CH₃), 8.59(2H, s, -NH quinazoline ring), 5.51(1H, s, =CH), 2.05(2H, t, CH₂, quinazoline ring), 1.39(2H, t, CH₂, quinazoline ring).

4-(1-(3-chlorobenzo[b]thiophene-2-carbonyl)-1H-indol-3-yl) 7, 7-dimethyl-3, 4, 7, 8-tetrahydroquinazoline-2, 5(1H, 6H) dione (3d)

IR (KBr) cm^{-1} : 3053(C-H str., Aromatic ring), 2958(C-H str., -CH₃ group), 3255(N-H str.), 687(C-S-C str., Benzo[b] thiophene ring), 769(C-Cl str.), 1680(C=O str.), 3380(OH str.); ^1H NMR (DMSO-d_6) δ : 6.99-8.11(7H, m, Ar-H), 1.26(6H, s, Ar-CH₃), 8.49(2H, s, -NH quinazoline ring), 5.56(1H, s, =CH), 5.79(1H, s, O-H), 1.99(2H, t, CH₂, quinazoline ring), 1.33(2H, t, CH₂, quinazoline ring).

Antimicrobial Activity

The newly synthesized compounds were screened for their antibacterial activity against gram positive bacteria *S. aureus* and gram negative *E. coli*, *Pseudomonas aeruginosa* and for antifungal activity against *Candida albicans*, *A. niger*, and *A. clavatus*. Antimicrobial activity was carried out by serial broth dilution method. The standard strains used for the antimicrobial activity was procured from Institute of Microbial Technology, Chandigarh. The compounds (3a-d) were screened for their antimicrobial activity at different concentrations of 1000, 500, 200, 100, 50 $\mu\text{g/ml}$

as shown in (Table 2). The results were recorded in the form of primary and secondary screening. The synthesized compounds were diluted at 1000µg/ml concentration, as a stock solution. The drugs which were found to be active in primary screening were similarly diluted to obtain 100, 50µg/ml concentrations. 10µg/ml suspensions were further inoculated on appropriate media and growth was noted after 24 and 48 h. The lowest concentration, which showed no growth after spot subculture was considered as MIC for each drug. The highest dilution showing at least 99% inhibition is taken as (MIC). The test mixture should contain 10⁸ cells/ml. The standard drug used in this study was 'Ampicillin' for evaluating antibacterial activity which showed (50, 100, and 50µg/ml) MIC against *E. coli*, *P. aeruginosa* and *S. aureus* respectively. For antifungal activity 'Griseofulvin' was used as a standard drug, which showed (100, 100, and 100µg/ml) MIC against *C. albicans*, *A. niger*, and *A. clavatus*, respectively.

RESULTS

Physical and analytical data of the synthesized compounds are given in Table 1. Table 2 describes results for antibacterial and antifungal activities. Synthesized compounds have been screened against three bacterial strains viz. *E. coli*, *P. aeruginosa*, *S. aureus* and three fungal strains viz. *C. albicans*, *A. niger*, *A. clavatus*. Compound **3c** is considered to be good active against *P. aeruginosa*. For the antifungal activity compounds **3c** is considered as good active against *A. niger* and *A. clavatus*.

Table 1: Physical and analytical data of synthesized compounds

Compound No.	Molecular formula	Molecular weight	Melting point (°C)	Yield (%)	Found/ Calculated (%) N
2a	C ₁₈ H ₁₀ ClNO ₂ S	339.79	276	81	4.10/4.12
2b	C ₁₈ H ₉ Cl ₂ NO ₂ S	374.24	152	79	3.72/3.74
2c	C ₁₈ H ₉ ClFNO ₂ S	357.78	285	80	3.89/3.91
2d	C ₁₈ H ₁₀ ClNO ₃ S	355.79	162	72	3.91/3.94
3a	C ₂₇ H ₂₂ ClN ₂ O ₃ S	503.99	480	73	8.32/8.34
3b	C ₂₇ H ₂₁ Cl ₂ N ₃ O ₃ S	538.44	336	68	7.78/7.80
3c	C ₂₇ H ₂₁ ClFN ₃ O ₃ S	521.99	484	71	8.03/8.05
3d	C ₂₇ H ₂₂ ClFN ₃ O ₄ S	519.99	342	70	8.02/8.08

Table 2: Biological activity of synthesized compounds (4a-d)

S. No.	Antibacterial Strains			Antifungal Strains		
	Gram negative		Gram positive	<i>C. albicans</i>	<i>A. niger</i>	<i>A. clavatus</i>
	<i>E. coli</i>	<i>P. aeruginosa</i>	<i>S. Aureus</i>			
3a	500	250	250	500	500	250
3b	250	250	100	500	250	500
3c	100	100	250	250	100	100
3d	100	250	100	100	100	250
S.D.*	50	100	50	100	100	100

S.D= Ampicillin for antibacterial drug; S.D= Griseofulvin for antifungal drug

ACKNOWLEDGEMENT

The authors are thankful to the Head, Department of Chemistry, M. L. Sukhadia University, Udaipur (Rajasthan) for providing laboratory facilities.

REFERENCES

1. Singh K, Sharma VK. Synthesis and antimicrobial study of (Z)-5-(substituted benzylidene)-2-(benzo[b]thiophene-2-

DISCUSSION

The NH proton of indol ring was replaced by various substituted 3-chlorobenzo[b]thiophene-2-carbonyl (**1a-d**) in ethanol using pyridine as a catalyst afforded (**2a-d**), were confirmed by a new band at 752 cm⁻¹ of C-Cl stretching and band at 687 due to C-S-C Stretching. Final compounds (**3a-d**) were synthesized using the Biginelli reaction, a multi component coupling of an equivalent moles of 5,5dimethylcyclohexane-1,3-dione, 1-[(3-chloro-1-benzothiophen-2-yl) carbonyl]-1H-indole-3-carbaldehyde (**2a-d**) and urea to furnished bicyclic compounds 4-(1-(3-chlorobenzo[b]thiophene-2-carbonyl)1H-indol-3-yl)-7,7-dimethyl-3,4,7,8-tetrahydroquinazoline-2,5(1H,6H)dione (**3a-d**). Structures of these compounds were confirmed by the presence of N-H stretching band at 3258 cm⁻¹. The 1H NMR spectral also confirms the structure of (**3a-d**), exhibited a singlet at δ 8.56 for N-H of quinazoline ring and a new singlet for =C-H proton of quinazoline ring at δ 5.56.

For antibacterial activity, compound **3c** is considered to be good active against *P. aeruginosa*, while for *S. aureus* compounds **3b**, **3d** and for *E. coli* compound **3c**, **3d** are moderately active. For the antifungal activity compound **3c** is considered as good active against *A. niger* and *A. clavatus* while compound **3d** against *C. Albicans* and *A. niger*. The discussion and comparison of antibacterial and antifungal activities have been compared with Ampicillin and Griseofulvin respectively.

carbonyl)-3-phenyl-1, 2-dihydro-1, 2, 4-triazin-6(5H)-one. Asian Journal of Biochemical and Pharmaceutical Research 2014; 4(3): 179-186, 8.

2. Dang H, Chorell E, Uvell H, Pinkner JS, Hultgren SJ, Almqvist F. Syntheses and biological evaluation of 2-amino-3-acyl-tetrahydrobenzothiophene derivatives; antibacterial agents with antivirulence activity. Organic & Biomolecular Chemistry 2014; 12(12): 1942-1956.

3. Naganagowda G, Tham Yongkit P, Petsom A. Synthesis and antimicrobial activities of benzothiophene derivatives. *Journal of the Chilean Chemical Society* 2012; 57(1): 1043-1047.
4. Vibhutamath GB, Mruthyunjayaswamy BHM. Synthesis and antimicrobial activity of some new N'-(3-chloro-6-substituted benzo[b]thiophene-2-carbonyl) substituted 2-oxo-2H-pyrano [2, 3-b] quinoline-3-carbohydrazides. *Organic Chemistry: An Indian Journal* 2011; 7(5): 325-331.
5. Shedid SAM, Hassan HM, Kora FA, El-Eisawy RM. Synthesis and antibacterial activity of some new amino acid derivatives of 3-chloro-6-(methyl)benzo[b]thiophene-2-carboxylic acid. *Journal of Chemical and Pharmaceutical Research* 2011; 3(1): 388-394.
6. Isloor AM, Kalluraya B, Sridhar PK. Synthesis, characterization and biological activities of some new benzo[b]thiophene derivatives. *European journal of medicinal chemistry* 2010; 45(2): 825-30.
7. Guruprasad BV, Mruthyunjayaswamy BHM. Synthesis and antimicrobial activity of some new 3-chloro-6-substituted-N-(substituted 2H-[1,3]oxazino[6,5-b]quinolin-3-(4H,5aH,9aH)-yl)benzo[b]thiophene-2-carboxamides. *Indian Journal of Chemistry, Section B: Organic Chemistry Including Medicinal Chemistry* 2012; 51B(3): 514520.
8. Tirilapur VK, Swamy KM, Prasad YR. Synthesis and antimicrobial studies of substituted benzo[b]thiophenes and their derivatives. *International Journal of Chemical Sciences* 2008; 6(4): 2008-2015.
9. Pinto E, Queiroz Maria-Joao RP, Vale-Silva LA, Oliveira JF, Begouin A, Begouin Jeanne-M, Kirsch G. Antifungal activity of synthetic di(hetero)arylamines based on the benzo[b]thiophene moiety. *Bioorganic & Medicinal Chemistry* 2008; 16(17): 8172-8177.
10. Yadav PP, Gupta P, Chaturvedi AK, Shukla PK, Maurya R. Synthesis of 4-hydroxy-1-methylindole and benzo[b]thiophen-4-ol based unnatural flavonoids as new class of antimicrobial agents. *Bioorganic & Medicinal Chemistry* 2005; 13(5): 1497-1505.
11. Tanaka J, Yuasa T, Kawakami Y, Terajima K, Ishizuka Y. Preparation of antifungal triazole-containing 2,3-dihydrobenzo[b]thiophene derivative. *Jpn. Kokai Tokkyo Koho*. 1996; JP 08253475 A 19961001.
12. Drouhet E, Dupont B. In vitro antifungal activity of sertaconazole. *Arzneimittel-Forschung*. 1992; 42(5A): 705-10.
13. Boyd JE, Somerville RG. Antiviral activity of some related benzo(b) thiophene derivatives. I. Initial screening tests. *Archiv fuer die Gesamte Virusforschung*. 1974; 45(3): 249-53.
14. Shimizu S, Hosoya T, Murohashi M, Yoshida S. Benzothiophene compound, alternative autophagy-inducing agent and anticancer drug including benzothiophene compound as active ingredient, and method for screening for compounds having anticancer activity. *PCT Int. Appl.* 2013; WO 2013118842 A1 20130815.
15. Penthala NR, Sonar VN, Horn J, Leggas M, Yadlapalli Jai Shankar KB, Crooks PA. Synthesis and evaluation of a series of benzothiophene acrylonitrile analogs as anticancer agents. *Med Chem Comm.* 2013; 4(7): 1073-1078.
16. Kagatani S, Odanaka J, Kano H. DNA methyltransferase inhibitors and antitumor agents containing benzo[b]thiophene derivatives. *Jpn. Kokai Tokkyo Koho*. 2007; JP 2007153792 A 20070621.
17. Li Z, Liu K. Preparation of benzo[b]thiophene derivatives as anti-inflammatory agents. *Faming Zhuanli Shenqing*. 2013; CN 103058988 A 20130424.
18. Lakshmesha CS, Venugopala Reddy KR, Mohan S. Synthesis and microwave assisted transformation of ethyl-4,5,6,7-tetrahydro benzo (b) thiophene-2-amino-3-carboxylate into potential anti-inflammatory agents- novel mono and diacids. *Archives of Applied Science Research* 2011; 3(5): 556-561.
19. Gouda, TS, Jagadish, Salahuddin, M Shantakumar SM. Synthesis and antiinflammatory activity of some novel benzo(b)thiophene substituted thiazolidinones. *Asian Journal of Chemistry* 2009; 21(7): 5171-5178.
20. Radwan Mohamed AA, Shehab MA, El-Shenawy SM. Synthesis and biological evaluation of 5-substituted benzo[b]thiophene derivatives as anti-inflammatory agents. *Monatshefte fuer Chemie*. 2009; 140(4): 445-450.
21. Fakhr IMI, Radwan MAA, El-Batran S, Abd El-Salam Omar ME, El-Shenawy SM. Synthesis and pharmacological evaluation of 2-substituted benzo[b]thiophenes as anti-inflammatory and analgesic agents. *European Journal of Medicinal Chemistry* 2009; 44(4): 1718-1725.
22. Svoboda J, Stadler M, Jandera A, Panajotova V, Kuchar M. Synthesis and biological evaluation of new antiinflammatory 1-benzothiophene-2-carboxanilides. *Collection of Czechoslovak Chemical Communications* 2000; 65(7): 1082-1092.
23. Yoshid, Y. Preparation of benzo[b]thiophene-2-carboxylic acids as intermediates for anti-inflammatory and antiallergy agents. *Jpn. Kokai Tokkyo Koho*. 1993; JP 05194484 A 19930803.
24. Banerjee T, Kapoor N, Surolia N, Surolia A. Benzothiophene carboxamide derivatives as novel antimalarials. *IUBMB Life*. 2011; 63(12): 1111-1115.
25. Bhatt R, Sharma S, Patidar AK, Rathore KK, Senwar RC, Dashora R, Mehta A. Synthesis and Antimicrobial evaluation of some imidazo imidazole Pyrimidines derivatives of benzo[b]thiophene. *Inter. J. of Pharmacy and Pharma. Sci. Res.* 2015; 5(1): 8-12.
26. Nagesh HK, Padmashali B, C. Sandeep, T.C.M. Yuvaraj, M.B. Siddesh, S. M. Mallikarjuna. Synthesis and Antimicrobial Activity of Benzothiophene Substituted Coumarins, Pyrimidines and Pyrazole as New Scaffold. *Int. J. Pharm. Sci. Rev. Res.* 2014; 28(2): 6-10.
27. Vivekanand D, Biradarand BHM, Mruthyunjayaswamy. Synthesis Characterization and Antimicrobial Activity Studies of Some Transition Metal Complexes Derived from 3-Chloro-N-[(1E)-(2-hydroxy phenyl)methylene]-6methoxy-1-benzothiophene-2-carbohydrazide. *Hindawi Publishing Corporation. The Scientific World Journal* 2013; Article ID 451629: 13.
28. Singh A, Singh SP, Amalraj A, Gokulan PD, Porwal P. Synthesis, Characterization and Biological Evaluation of 2-amino-3-(n-cyclohexyl carboxamido)-4,5,6,7-tetrahydrobenzo (b) Thiophine Derivatives. *Inter. J. of Pharma. & Bio. Archives*. 2011; 2(1): 534-540.
29. Vasoya SL, Paghdar DJ, Chovatia PT, Joshi HS. Synthesis of some New Thiosemicarbazide and 1,3,4Thiadiazole Heterocycles Bearing Benzo[b]Thiophene Nucleus as a Potent Antitubercular and Antimicrobial Agents. *J. of Sci., Islamic Republic of Iran* 2001; 16(1): 33-36.
30. Thakar KM, Paghdar DJ, Chovatia PT and Joshi HS. Synthesis of thiourea derivatives bearing the benzo[b]thiophene nucleus as potential antimicrobial agents. *J. Serb. Chem. Soc.* 2005; 70 (6): 807-815.
31. Kumara THS, Mahadevan KM, Harishkumar HN, Padmashali B, Naganagowda G. Synthesis of benzo[b]thiophene substituted carbamates, ureas, semicarbazides, and pyrazoles and their antimicrobial and analgesic activity. *Phosphorus, Sulfur and Silicon and the Related Elements* 2009; 184(7):1866-1879.

Source of Support: Nil, Conflict of Interest: None declared.