



Pharmacognostical Evaluation of Roots of *Simmondsia chinensis* Schneider

Surendra Kumar Sharma*, Ajay Pal Singh

Department of Pharmaceutical Sciences, Guru Jambheshwar University of Science and Technology, Hisar-125 001, Haryana, India

ABSTRACT

Simmondsia chinensis Schneider commonly known as jojoba is a semi-arid evergreen shrub. Jojoba is unique among plants in the fact that its products (seeds) contain about 50 percent by weight oil. Jojoba liquid wax generally used in folk remedies for renal colic, sunburn, chaffed skin, hair loss, headache, wounds and sore throat. However, there are no reports on the roots of this plant. The present investigation deals with the pharmacognostical studies of the roots of the said plant. Pharmacognostical studies include microscopic, physico-chemical constants (ash & extractive values), fluorescence analysis and preliminary phytochemical evaluation.

Keywords: *Simmondsia chinensis*, physico-chemical parameters, phytochemical screening, fluorescence analysis.

INTRODUCTION

Simmondsia chinensis Schneider commonly known as jojoba is a semi-arid evergreen shrub. [1] It grows wild in desert of south-western United States and north-western Mexico. However, this plant is cultivated in Australia [2], Brazil, Argentina, Middle East countries [3] and India. [4] Jojoba is unique among plants in the fact that its products (seeds) contains about 50 percent by weight oil, which is more than amount in soyabean and somewhat more than in most oil seed crops. Interest in jojoba oil stems from its unusual properties that differ from all seed oils. The complete absence of glycerin makes it a liquid wax (JLW), not fat. [5] Jojoba liquid wax contains a natural anti-oxidant postulated to be an allylic derivative of hydroxytoluene. [6] Jojoba liquid wax was used in folk remedies for renal colic, sunburn, chaffed skin, hair loss, headache, wounds and sore throat. [7] Animal studies demonstrate that Jojoba liquid wax can be classified as a non-toxic substance. [8] Jojoba liquid wax has shown moderate digestibility that increases when the wax is mixed with other oils. In addition, Jojoba liquid wax possesses moderate absorption when applied topically. [9] In human studies, sulphurized jojoba liquid wax was effective in the treatment of acne while the unmodified wax was used

for treatment of psoriasis. [10] Furthermore, dermatological research suggests that Jojoba liquid wax may help to reduce inflammation. [11] A lot of works as suggested by literature survey had been carried out on seeds and seed oil but, there are no reports on roots of *Simmondsia chinensis*. The present study was therefore undertaken to investigate the pharmacognostical characters, fluorescence analysis and phytochemical analysis of the roots of this plant.

MATERIAL AND METHODS

Plant material

The plant material was collected from Kutch district, Gujarat, July 2009. The plant material was identified and authenticated by the Dr. H. B. Singh, Head, Raw Materials Herbarium & Museum, NISCAIR, New Delhi, Ref. No. NISCAIR/ RHMD/Consult/-2009-10/1204/08. The dried roots were ground to a coarse powder and subjected to physico-chemical analytical parameters like ash values, extractive values and phytochemical screening.

Chemicals and instruments

Rotary microtome was used to take sections of the tubers. Compound microscope, glass slides, cover slips, watch glass and other common glassware were used in this experiment. Photographs were taken with using Zeiss Primo Star Microscopic Unit. Various solvents used mainly ethanol (95%), petroleum ether, chloroform and reagents used for staining different sections were procured from S D Fine chemicals, Mumbai, India.

Morphology

A systematic examination of the shape, size, surface, texture, taste and odour of the roots of *Simmondsia chinensis* was

*Corresponding author: Dr. Surendra Kumar Sharma, Professor and Chairman, Department of Pharmaceutical Sciences, & Dean, Faculty of Medical Sciences, Guru Jambheshwar University of Science and Technology, Hisar-125 001, Haryana, India; Tel.: +91-01662-263580; Fax: +91-01662-276025; E-mail: prof.sharmask@gmail.com, ajaypharma@gmail.com

carried out. The external features of root were observed under dissecting microscope.

Microscopy

The histological examination of the plant drug helps in identifying the shapes, size and position of different cells and tissues. The anatomical study helps in distinction of the drug from adulterants. [12] Material collected (were fixed in formalin – acetic acid – alcohol (10.5: 50: 35). The fixed material was preserved in 70% alcohol, washed thoroughly in 70% alcohol, dehydrated through xylene – alcohol series and embedded in paraffin. [13] Serial transverse sections were cut at 7-8 μ m on a rotary microtome (Weswox Optik). Hapt adhesive was used for affixing the paraffin ribbons to the slide. The sections were stained in phloroglycerol. Slides were cleared in xylol and mounted in DPX mountant. Microscopic descriptions of tissues are supplemented with micrographs wherever necessary. Photographs of different magnifications were taken with Cannon Zeiss Primo Star Microscopic unit. [14]

Powder microscopic study

The shade dried plant materials were ground with a wood grinder and sifted through 40 mesh sieves. To study the ingredients of powder, a pinch of powder was bleached with 5% chloral hydrate and taken on slide and mounted with phloroglucinol, HCl and glycerine. The slide was observed under microscope.

Colour reactions

The powdered drug materials were treated with different chemical reagents so as to aid in detection of chemical constituents under ordinary day light by standard methods. [15] To study the behaviour of drugs a pinch of each drug was treated with different chemical reagents *viz* 1N HCl, 1N NaOH, acetic acid, 5% ferric chloride, picric acid, 1N HNO₃, 5% iodine and 1N HNO₃ followed by ammonia solution and colours were observed. [16]

Fluorescence behaviour

To study the fluorescence nature of powder, a pinch of powder after bleaching with 5% chloral hydrate was treated with different chemical reagents *viz*. 1N HCl, 1N NaOH, 50% HNO₃, 50% H₂SO₄, methanol, acetic acid, picric acid, 1N NaOH in methanol, nitro-cellulose in amyl acetate, 1N NaOH in methanol and nitrocellulose in amyl acetate, 1N HCl followed by nitrocellulose in amyl acetate and 1N NaOH followed by nitrocellulose in amyl acetate and observed under UV light. [16-17]

Physico-chemical parameters

The determination of various physico-chemical parameters such as total ash, acid insoluble ash, water soluble ash, sulphated ash, water soluble extractive and alcohol soluble extractive were determined and estimated in percentage by using method as recommended by Indian Pharmacopoeia. [18] Successive soxhlet extractives of the drug were carried out with various solvents and weight, color/consistency of extractives were observed. [17] Loss on drying for drug was also determined. [19-20]

Preliminary phytochemical screening

The powdered roots extracts were subjected to qualitative phytochemical tests for alkaloids, glycosides, carbohydrates, sterols, phenolic compounds and tannins, flavonoids, saponins, proteins and free amino acids. [21-22]

RESULTS AND DISCUSSION

Macroscopic characters

It occurs as tap root system up to 15 feet. Externally it is blackish brown and internally reddish brown in colour. Surface is rough with striations. Odour is characteristic and taste is woody.




Fig. 1. Transverse section of *Simmondsia chinensis* root (Co-Cork; Cr-Cortex; Ph-Phloem; MR-Medullary rays; Ve-Vessel and XF-Xylem Fibres)




Fig. 2. Microscopic characteristics of powder of *Simmondsia chinensis* root Ve-Vessel, Co- Cork, F- Fibre and St-Starch

Table 1: Behavior of root powder with different chemical reagents

| S. No. | Treatment | Colour |
|--------|--|-----------------|
| 1. | Powder | Dark brown |
| 2. | Powder + 1 N HCl | Yellowish brown |
| 3. | Powder + 1N NaOH | Black |
| 4. | Powder + Acetic Acid | Brownish black |
| 5. | Powder + 5% Ferric chloride | Yellowish brown |
| 6. | Powder + Picric acid | Greenish brown |
| 7. | Powder + HNO ₃ + Ammonia solution | Yellowish brown |
| 8. | Powder + 5% Iodine | Brown |
| 9. | Powder + 1N HNO ₃ | Reddish brown |

Table 2: Fluorescent nature of *Simmondsia chinensis* root powder

| S. No. | Treatment | Visible | Observations | |
|--------|---|-----------------|-------------------|-----------------|
| | | | Short UV (254 nm) | Long UV (366nm) |
| 1. | Powder as such | Brown | Green | Dark brown |
| 2. | Powder + 1N HCl | Yellowish brown | Greenish black | Light brown |
| 3. | Powder + 1N NaOH | Black | Green | Dark brown |
| 4. | Powder + 50% HNO ₃ | Reddish brown | Black | Greenish black |
| 5. | Powder + 50% H ₂ SO ₄ | Brown | Greenish brown | Black |
| 6. | Powder + Methanol | Creamish brown | Brown | Dark brown |
| 7. | Powder + Acetic acid | Brownish black | Greenish brown | Black |
| 8. | Powder + Picric acid | Yellowish brown | Brownish green | Black |
| 9. | 5% Iodine | Brown | Dark brown | Black |
| 10. | 5% Ferric chloride | Yellowish brown | Yellowish black | Black |
| 11. | Ammonia | Brown | Greenish brown | Black |

Table 3: Physico-chemical parameters of roots powder of *Simmondsia chinensis*

| Physico-chemical parameters | Values [% (W/W) ± SD] |
|-----------------------------|-----------------------|
| Ash values | |
| Total ash | 9.06 ± 0.002 |
| Acid insoluble ash | 3.73 ± 0.004 |
| Water soluble ash | 6.15 ± 0.007 |
| Sulphated ash | 9.16 ± 0.003 |
| Extractive values | |
| Water soluble extractive | 8.01 ± 0.011 |
| Alcohol soluble extractive | 9.46 ± 0.008 |

Data are mean ± SD values (n = 3)

Table 4: Yield of extract by successive solvent extraction

| S. No. | Extract* | Values [% (W/W) ± SD] |
|--------|-----------------|-----------------------|
| 1. | Petroleum Ether | 02.13 ± 0.002 |
| 2. | Chloroform | 03.31 ± 0.005 |
| 3. | Methanol | 10.12 ± 0.010 |
| 4. | Water | 05.76 ± 0.013 |

*Extraction period is 24 hrs, Data are mean ± SD values (n = 3)

Table 5: Loss on drying

| Wt. of the crucible | Wt. of crucible+ Drug | Final wt. | %w/w |
|---------------------|-----------------------|-----------|-------|
| | | 37.21g | 6.98% |
| 35.61g | 37.33g | 37.20g | 7.56% |
| | | 37.20g | 7.56% |

Table 6: Fluorescence nature of different extracts of *Simmondsia chinensis* by visible and ultra-violet (UV) radiations

| S. No. | Extract | Observations | | |
|--------|-----------------|--------------|-------------------|-----------------|
| | | Visible | Short UV (254 nm) | Long UV (366nm) |
| 1. | Petroleum ether | Yellow Brown | Brown | Black |
| 2. | Chloroform | Brown | Greenish Brown | Black |
| 3. | Ethanol | Brown | Dark Brown | Dark Brown |
| 4. | Water | Black Brown | Brown | Black |

Microscopic characters

Table 7: Preliminary phytochemical screening of roots extract of *Simmondsia chinensis*

| S. No. | Plant Constituents | Petroleum Ether Extract | Chloroform Extract | Ethanol Extract | Water Extract |
|--------|-----------------------|-------------------------|--------------------|-----------------|---------------|
| 1. | Alkaloids | - | - | - | - |
| 2. | Glycosides | - | - | - | - |
| 3. | Protein & Amino acids | - | + | + | + |
| 4. | Carbohydrates | - | - | + | + |
| 5. | Tannins & Phenolics | - | - | + | + |
| 6. | Flavonoids | - | + | + | + |
| 7. | Steroids | + | + | - | - |
| 8. | Saponins | - | - | - | - |
| 9. | Fixed oil | + | - | - | - |

+ : Positive; - : Negative

Transverse section of the roots shows the following characteristics:

Cork is composed of 3-5 layers of tangentially elongated cells filled with brown matter. Phellogen is indistinct but seen as narrow layers of cells. Phelloderm 5-7 layers immediately below the phellogen. Cortex is composed of several layers of polyhedral parenchymatous cells. Phloem present as several thin patches around the well developed xylem. Secondary xylem is transverse by well develop medullary rays. Xylem consists of vessels, wood parenchyma. The vessel appears rounded or polygonal. Xylem fibres appear as rounded and polygonal structure with thick lignified walls. Medullary rays runs radially from the centre of the cortex through the phloem. Rays in the xylem region are lignified, pitted and are 1-3 cells wide although uniserate are prominent. In the phloem regions the ray cells are not lignified (Fig. 1).

Powder characters

Microscopic examination of powder shows the presence of cork cells, starch, vessels and fibres (Fig. 2). The behavior of roots powder with different chemical reagents, are shown in (Table 1).

The fluorescence analysis of powdered drug in day light, short UV and long UV were examined by reported method. The observations are given in (Table 2).

Physico-chemical parameters

The physico-chemical parameters are important for identifying adulterants and improper handling of drugs. The (Table 3) reveals the result of physico-chemical parameters of powdered drug, carried out by using standard procedures. viz. Ash values used to determine quality and purity of drug; the extractive values are useful to evaluate the chemical constituents present in crude drugs and also help in estimation of specific constituents soluble in particular solvent. The result of extractive values of powdered drug in different solvent obtained by successive extraction is given in (Table 4). Loss on drying is shown in (Table 5). The (Table 6) reveals the behavior analysis of different solvent extract of *Simmondsia chinensis* root, under visible light, short and long UV.

All the extracts obtained by successive extraction were subjected to qualitative chemical tests and the results are shown in (Table 7). Such preliminary phytochemical screening is helpful in prediction of the nature of drugs and also useful for detection of different constituents in different polarity solvent.

The present investigation including the morphoanatomical characters, physico-chemical values and pharmacognostical

studies will serve as standard reference for identification and distinguishing *Simmondsia chinensis* root from its substitute and adulterants. This report would assist in the identification of the crude drug in future.

REFERENCES

1. Harry-O'Kura RE, Mohamed A, Abott TP. Synthesis and characterization of tetrahydroxyjojoba wax and ferulates of jojoba oil. *Industrial Crops and Products* 2005; 22: 125-133.
2. Davidson S. Jojoba: cautious optimism. *Rural Res.* 1983; 119: 21-25.
3. Borlaug N, Baldwin AR, Estefan R, Harris m, Plucknett DL. Jojoba: new crop for arid lands. New raw material for industry. National Academy Press, Washington, 1985, pp. 6, 13.
4. Kaul S, Goyal HB, Bhatnagar AK, Gupta AK. Effect of ageing on quality of jojoba oil from Indian locations. *Industrial Crops and Products* 2009; 29: 102-107.
5. Abu-Arabi MK, Allawzi MA, Al-Zoubi HS, Tamimi A. Extraction of jojoba oil by pressing and leaching. *Chem. Eng. J.* 2000; 76: 61-65.
6. Kampf A, Gringberg S, Galuun A. Oxidative stability of jojoba wax. *J. Am. Oil Chem. Soc.* 1986; 63:246-248.
7. Yaron A. Metabolism and physiological effects of Jojoba oil. In: Wisniak J, editor. *The chemistry and technology of jojoba oil.* American Oil Chemists' Society Press, Champaign, 1987, pp. 251-65.
8. Taguchi N, Kunitomo T. Toxicity studies on jojoba oil for cosmetic uses. *Cosmet. Toilet* 1977; 92: 53-62.
9. Heise C, Decombaz J, Anantharaman K. Energy value of jojoba oil for the growing rat. *Int. J. Vitam. Nutr. Res.* 1982; 52: 216-221.
10. Mosovich B. Treatment of acne and psoriasis. In: Wisniak J, Zabicky J, editors. *Proceedings of the Sixth International Conference on Jojoba and its Uses.* Beer Sheva, Israel: University of the Negev; 1985. pp. 393-7.
11. Borlaug N, Baldwin AR, Estefan R, Harris M, Plucknett DL. Jojoba: new crop for arid lands. New raw material for industry. National Academy Press, Washington DC, 1985. pp. 53.
12. Ali M. *Textbook of Pharmacognosy.* Edn 1, CBS Publishers and Distributors, New Delhi, 1998, pp. 24-28.
13. Panday BP. *Modern Practical Botany.* Edn 1, S. Chand Publishers, New Delhi, 1980, pp. 1-6.
14. Johansen DA. *Plant Microtechnique.* Mc Graw Hill Book Co, New York, 1940, p. 523.
15. Chase CR, Pratt RJ. Fluorescence analysis of powdered vegetable drugs with particular reference to development of a system of identification. *J. Am. Pharmacol. Assoc.* 1949; 38: 324-331.
16. Kokoski CJ, Kokoski RJ, Salma FJ. Fluorescence of powdered vegetable drugs under UV radiation. *J. Am. Pharmacol. Assoc.* 1958; 47: 715-717.
17. Mukherjee PK. *Quality Control of Herbal Drug.* Edn 1, Business Horizons Pharmaceutical Publishers, New Delhi, 2010, pp. 184-191.
18. Anonymous: *Indian Pharmacopoeia*, Ministry of Health and Family Welfare, Government of India, Controller of Publications, New Delhi, 1996, pp. A48.
19. World Health Organization (WHO): *Quality Control Methods for Medicinal Plant Materials*, Office of Publications, England, 1998, pp. 8.
20. Yadav N, Vasudeva N, Sharma SK, Singh S. Pharmacognostical and phytochemical studies on *Chenopodium album* Linn. root. *Hamdard Medicus* 2007; 50: 95-102.
21. Harborne JB. *Phytochemical Methods.* Edn 2, Chapman and Hall, London, 1973, pp. 5-6.
22. Kokate CK. *Practical Pharmacognosy.* Edn 4, Vallabh Prakashan, Pune, 1996 pp. 107-109.