

International Journal of Pharmaceutical Sciences and Drug Research

2016; 8(4): 240-243



Research Article

ISSN: 0975-248X
CODEN (USA): IJPSPP

Isolation and Identification of Steroid and Flavonoid Glycosides from the Flowers of *Allium gramineum*

Lasha Mskhiladze^{1*}, David Chincharadze¹, Alexis St-Gelais², Vakhtang Mshvildadze², André Pichette², Allison Ledoux³, Monique Tits³, Michel Frédérich³

¹Department of Pharmacognosy and Botany, Faculty of Pharmacy, Tbilisi State Medical University, 33, Vazha Pshavela Ave., Tbilisi, 0177, Georgia

²Laboratory LASEVE, Department of Sciences Fondamentales, University of Québec in Chicoutimi, 555 Boul. of University, Chicoutimi, Québec, Canada

³Laboratory of Pharmacognosy, Department of Pharmacy, Centre Interdisciplinaire de Recherche du Médicament-CIRM, University of Liège, CHU-B36, B-4000 Liège, Belgium

ABSTRACT

The isolation and identification of β -sitosterol 3-O- β -glucopyranoside, quercetin 3-O- β -glucopyranoside, isorhamnetin 3,4'-di-O- β -glucopyranoside and isorhamnetin 3,7-di-O- β -glucopyranoside from the flowers of *Allium gramineum* that grows in Georgia. The structures of isolated compounds have been determined by ¹H and ¹³C nuclear magnetic resonance (NMR).

Keywords: *Allium gramineum*, Alliaceae, Steroid glycoside, Flavonoid glycoside, Structure determination..

INTRODUCTION

Over the past 10 years, researchers have focused increasing attention on extraction, identification, fractionation and isolation of phenolic compounds in plants. They have great abundance in our diet, and they have a probable role in the prevention of various diseases associated with oxidative stress, such as cancer as well as cardiovascular and neurodegenerative diseases. [1-4] Flavonoids represent a large group of phenolic compounds found in plants, these compounds demonstrable human health related benefits including

antioxidant status, antiviral, anti-inflammatory and anti-cancer properties. [5-10]

Allium species are rich sources of steroid and flavonoid glycosides, among which quercetin and isorhamnetin glucosides are the major components. They are known for their antioxidant and free radical scavenging potentialities, their antiallergic, anti-inflammatory and cytotoxic activities. [11-14] A study using the prostate cancer cell line LNCaP (an androgen-dependent tumor) showed that β -sitosterol-3-O- β -D-glucoside decreased cancer cell growth by 24% and induced apoptosis (programmed cell death) fourfold, with little side effect. Also, β -sitosterol-3-O- β -D-glucoside characterized by antioxidant and free radical scavenging activities. [15-16] *Allium gramineum* (family Alliaceae) is an indigenous *Allium* species from Caucasus and, along with other species, is widely used in Georgian traditional medicine as an antifungal, antiseptic and antibacterial

*Corresponding author: Mr. Lasha Mskhiladze, Assistant Professor, Department of Pharmacognosy and Botany, Faculty of Pharmacy, Tbilisi State Medical University, 33, Vazha Pshavela Ave., Tbilisi, 0177, Georgia; Tel.: +995 599409555; Fax: +995 32 2340756; E-mail: lashamskhiladze@gmail.com
Received: 23 July, 2016; Accepted: 27 July, 2016

remedy. [17] Phytochemical and biological studies of *Allium* species growing in Georgia have been previously reported and showed promising results. [18-21] This prompted use to continue in-depth the phytochemical study of the flowers of *Allium gramineum* growing in Georgia. This paper describes the isolation and identification of β -sitosterol 3-O- β -glucopyranoside, quercetin 3-O- β -glucopyranoside, isorhamnetin 3, 4'-di-O- β -glucopyranoside and isorhamnetin 3, 7-di-O- β -glucopyranoside from the flowers of *Allium gramineum*. The structures of isolated compounds described for the first time in *Allium gramineum*, were determined by ^1H and ^{13}C nuclear magnetic resonance (NMR).

MATERIALS AND METHODS

General experimental

For column chromatography, silica gel 60 (40-63 μm , Merck) and Diaion HP-20 resin (Mitsubishi) were used. TLC analysis were performed on Silica gel 60 F254 plates (Merck) and eluted with CH_2Cl_2 -MeOH- H_2O (55:12:1, v/v/v). Spots were detected by spraying the plates with vanillin-sulphuric acid (in EtOH) reagent, followed by heating at 110°C . Preparative HPLC was a Varian PrepStar 218 coupled with a DAD (diode-array detector) detector set at 350 nm and equipped with a fraction collector (440LC) on a C-18 column, using a binary solvent system with a flow rate of 30 mL/min: solvent A: trifluoroacetic acid 0.05% in water and solvent B: acetonitrile. The Purity of isolated compounds was estimated on Agilent 1100 HPLC with DAD (diode-array detector) detection (working wavelength : 350 nm) using Hypersil ODS (C-18) columns (58 μm , 4.6 $\times$ 250 mm) with the same binary solvent system as described above, injection volume 10 μL with a flow rate of 1 mL/min. The isolated compounds were identified by NMR. ^1H and ^{13}C NMR spectra were recorded on a Bruker Avance DRX 500 instrument equipped with a cryoprobe. ^1H and ^{13}C NMR chemical shifts in ppm were referenced to the residual solvent (DMSO) signals or to TMS as internal standard (^1H at 500 MHz and ^{13}C at 125 MHz). 2D experiments were performed using standard Bruker microprograms.

Plant material

The whole plants of *Allium gramineum* were collected in the Didgori region of Georgia in June 2014, and identified by Dr. J. Aneli (Iovel Kutateladze Institute of Farmacochemistry, Tbilisi State Medical University). A voucher specimen (Flowers #133,134 TBPB) was deposited at the herbarium of Iovel Kutateladze Institute of Farmacochemistry, Tbilisi State Medical University.

Extraction and isolation

Dried and powdered flowers of *Allium gramineum* (100 g) were extracted twice with hot EtOH- H_2O (8:2 v/v 3 $\times$ 500 ml) to give the total extract (Ext.-1) After evaporation of the solvent, the residue Ext.-1 (30.8 g) was chromatographed over Diaion HP-20, using H_2O -MeOH as eluent in gradient conditions (100:0; 50:50; 0:100 v/v each 2 L) and finally EtOAc (1 L) to give four enriched fractions (F-1; F-2; F-3; F-4). After removing the solvents, the fraction F-1 (1.3 g) which was collected in H_2O 100% was subjected to silica gel CC and eluted with CH_2Cl_2 -MeOH- H_2O (55:12:1, v/v/v) to give: compound 1 (23 mg). The fraction F-2 (1.7 g) which was collected in H_2O -MeOH 50% was purified by Preparative HPLC coupled with a DAD detector set at 350 nm, using a binary solvent system: A: trifluoroacetic acid 0.05% in water and solvent B: acetonitrile (0 to 1 min 0% B, 100%A; 1 to 45 min 3%B, 97%A; 45 to 55 min 40%B, 60%A, 55 to 56 min 60%B, 40%A, 56 to 66 min 60%B, 40%A, 66 to 67 min 0%B, 100%A and 67 to 82 min 0%B, 100%A) to give: compound 2 (21 mg) compound 3 (16 mg) compound 4 (14 mg).

RESULTS AND DISCUSSION

A 80% EtOH extract of the flowers of *Allium gramineum* was subjected to successive fractionations to provide one steroid and flavonoid glycosides. Comparison of the ^1H - and ^{13}C -NMR spectral data with those reported in the literature led to the identification of the structures of compounds 1, 2, 3, 4 (Figure 1), described for the first time in *Allium gramineum*.

Compound 1 was identified as β -sitosterol 3-O- β -glucopyranoside by comparison with literature. [22] Compound 2 was obtained as an amorphous yellow powder. The APT spectrum indicated the presence of one methylene, ten methines and ten quaternary carbons, including one carbonyl.




Fig. 1: Chemical structures of compound 1, 2, 3, 4.

The COSY spectrum indicated that aromatic methines H-5' and H-6' were vicinal. The HMBC spectrum suggested that the aromatic ring was completed with oxygenated C-4' and H-2' in *meta* of the doublet doublet H-6'; and C-1' and oxygenated C-3' in *meta* of the doublet H-5'. Proton H-6' also exhibited a HMBC correlation with the downshifted C-2, suggesting it as the attachment point of the aromatic ring. Furthermore, another aromatic system was determined from *meta* coupling doublets H-6 and H-8. Both protons showed HMBC correlations with oxygenated C-7, and quaternary C-10, while H-6 correlated with oxygenated C-5 and H-8 with oxygenated C-9. These information, along with the presence of a relatively upshifted ketone, suggested a flavonol moiety with oxygenated groups in positions 3, 5, 7, 3' and 4', corresponding to a quercetin scaffold. Of these positions, only C-3 showed a HMBC correlation with H-1'', the others being free phenol groups. Along with the downshifted H-1'' signal in the anomeric region, a hexose moiety was inferred from the presence of the methylene and four remaining methines in the glycosidic region of the 1D NMR spectra. Proton H-1'' showed a HMBC correlation with C-5'', which suggested a pyranosyl conformation. Comparison with literature confirmed that 2 was quercetin 3-O- β -glucopyranoside, also known as isoquercetin. The large coupling constant of 7.6 Hz for proton H-1'' confirmed that the sugar moiety was the β anomer of glucose. [23]

Compound 3, isolated as an amorphous yellow powder, had an APT spectrum similar to that of 2, to the exception of a downshift of C-3' and the presence of one methyl, one more methylene and five additional methines. The methyl group chemical shift corresponded to a methoxy group, whose protons showed a HMBC correlation with C-3', indicating that the flavone was an isorhamnetin scaffold. The other additional signals were attributed to a second hexose moiety from their chemical shifts. As in 2, the anomeric proton H-1'' correlated with C-3, while the other anomeric signal H-1''' correlated with C-4'. Both H-1'' and H-1''' correlated with their 5 counterpart, indicating pyranoses, and exhibited a large coupling constant corresponding to their β anomers. [23] Furthermore, the chemical shifts of both hexoses were closely similar to those of the glucosyl of 2, suggesting that both were glucose residues. Thus, 3 were identified as isorhamnetin 3, 4'-di-O- β -glucopyranoside. Since no ^{13}C NMR reference could be found for this compound, data were compared with the spectrum of the equivalent quercetin diglucoside. [24] The two molecules showed a very similar NMR behavior except for positions 2', 3' and 4', which are slightly downshifted in compound 3 owing to the 3' methoxyl.

Compound 4, obtained as an amorphous yellow powder, showed an APT spectrum closely resembling that of 3. Examination of the HMBC correlations indicated that C-3 was again substituted with a

glucoside, but that the second hexose moiety was in position 7 of the flavone core. Both anomeric protons again exhibited a large coupling constant. Thus, the structure of 4 was determined as being isorhamnetin 3, 7-di-O- β -glucopyranoside. In absence of retrievable reference for the ^{13}C NMR spectrum of this flavonoid, the molecule was compared to its analogue quercetin 3, 7-di-O- β -glucopyranoside [25], with good agreement except for C-2' and C-3', near the methoxy substituent. Interestingly, compounds 2 and 3 have already been reported in *Allium cepa* and *A. ascalonicum*. [26-27]

Isorhamnetin 3, 4'-O- β -diglucopyranoside (3). ^1H NMR (500 MHz, DMSO-*d*₆) δ : 6.22 (d, *J* = 2.1, H-6), 6.47 (d, *J* = 2.1, H-8), 7.97 (d, *J* = 2.0, H-2'), 7.23 (d, *J* = 8.9, H-5'), 7.54 (dd, *J* = 8.9, 2.0, H-6'), 5.58 (d, *J* = 7.3, H-1''), 3.21 (m, H-2''), 3.10 (m, H-3'', H-4''), 3.23 (m, H-5''), 3.59 (d, *J* = 11.7, H-6''a), 3.39 (dd, *J* = 11.7, 5.7, H-6''b), 5.06 (d, *J* = 7.6, H-1'''), 3.29 (m, H-2''', H-5'''), 3.36 (m, H-3'''), 3.18 (m, H-4''') 3.67 (d, *J* = 11.7, H-6''a), 3.46 (dd, *J* = 11.7, 5.5, H-6''b), 3.84 (s, 3'-OCH₃); ^{13}C NMR (125 MHz, DMSO-*d*₆; APT) δ : 155.7 (C-2), 133.4 (C-3), 177.5 (C-4), 161.2 (C-5), 98.8 (C-6), 164.5 (C-7), 93.8 (C-8), 156.5 (C-9), 104.1 (C-10), 123.6 (C-1'), 113.5 (C-2'), 148.1 (C-3'), 148.5 (C-4'), 114.5 (C-5'), 121.4 (C-6'), 100.7 (C-1''), 74.3 (C-2''), 77.5 (C-3''), 69.8 (C-4''), 76.5 (C-5''), 60.6 (C-6''), 99.5 (C-1'''), 73.1 (C-2'''), 77.1 (C-3'''), 69.6 (C-4'''), 76.8 (C-5'''), 60.6 (C-6'''), 55.7 (3'-OCH₃).

Isorhamnetin 3, 7-O- β -diglucopyranoside (4). ^1H NMR (500 MHz, DMSO-*d*₆) δ : 6.45 (d, *J* = 2.1, H-6), 6.81 (d, *J* = 2.1, H-8), 7.95 (d, *J* = 2.0, H-2'), 6.93 (d, *J* = 8.5, H-5'), 7.53 (d, *J* = 8.5, 2.0, H-6'), 5.58 (d, *J* = 7.3, H-1''), 3.23 (m, H-2''), 3.10 (m, H-3'', H-4''), 3.27 (m, H-5'', H-2''', H-5'''), 3.58 (d, *J* = 11.2, H-6''a), 3.38 (dd, *J* = 11.2, 4.3, H-6''b), 5.08 (d, *J* = 7.3, H-1'''), 3.44 (m, H-3''', H-6''b), 3.15 (m, H-4'''), 3.69 (d, *J* = 9.5, H-6''a), 3.84 (s, 3'-OCH₃); ^{13}C NMR (125 MHz, DMSO-*d*₆; APT) δ : 156.9 (C-2), 133.3 (C-3), 177.6 (C-4), 160.9 (C-5), 99.4 (C-6), 162.9 (C-7), 94.6 (C-8), 156.0 (C-9), 105.7 (C-10), 120.9 (C-1'), 113.5 (C-2'), 146.9 (C-3'), 149.6 (C-4'), 115.2 (C-5'), 122.2 (C-6'), 100.7 (C-1''), 74.3 (C-2''), 77.5 (C-3''), 69.8 (C-4''), 76.5 (C-5''), 60.6 (C-6''), 99.7 (C-1'''), 73.1 (C-2'''), 77.2 (C-3'''), 69.6 (C-4'''), 76.4 (C-5'''), 60.6 (C-6'''), 55.7 (3'-OCH₃).

ACKNOWLEDGEMENTS

The authors wish to thank Dr. J. Aneli for identification of *A. gramineum*, Jean-Noël Wauters for technical help. This research has been partially supported by the program Shota Rustaveli National Science Foundation "President Research Grants for Young Scientists 2014" (Grant Agreement Nr. 52/38) and by the Belgian National Fund for Scientific Research (Grant Nr. T.0190.13).

REFERENCES

1. Boskou D. Sources of natural antioxidants. Trends Food Sci. Technol. 2006; 17: 505-512.
2. Maqsood S, Benjakul S, Abushelaibi A, Alam A. Phenolic Compounds and Plant Phenolic Extracts as Natural Antioxidants in Prevention of Lipid Oxidation in Seafood: A

- Detailed Review. *Comp. Rev. Food Science and Food Safety*. 2014; 13: 1125-1140.
3. Oscar D, Rangel H, Belen PV, Concepcion M, Aguilera G, Angel G. A Systematic Review of the Efficacy of Bioactive Compounds in Cardiovascular Disease: Phenolic Compounds. *Nutrients*. 2015; 7: 5177-5216.
4. Sytar O, Brestic M, Rai M, Shao HB. Plant phenolic compounds for food, pharmaceutical and cosmetics production. *Journal of Medicinal Plants Research*. 2012; 6 (13): 2526-2539.
5. Santhalakshmi R, Devaraj H, Niranjali DS. Quercetin Suppresses Twist to Induce Apoptosis in MCF-7 Breast Cancer Cells. *PLOS ONE*. 2015; 10 (10): 1-21.
6. Xu YJ, Zhang J, Liu S, Li C, Li W, Wang RM, Liu Y. Luteolin attenuate the D-galactose-induced renal damage by attenuation of oxidative stress and inflammation. *Natural Product Research*. 2015; 29: 1078-1082
7. Lu HY, Chen X, Sun B, Tong XH. Effects of luteolin on retinal oxidative stress and inflammation in diabetes. *RSC Advances*. 2014; 7: 4898-4904.
8. Jung H, Shin SY, Jung Y, Thao A, Lee Hk, Jung KY, Koh D, Cho SK, Lim Y. Quantitative Relationships Between the Cytotoxicity of Flavonoids on the Human Breast Cancer Stem-Like Cells MCF7-SC and Their Structural Properties. *Chem Biol Drug Des*. 2015; 86: 496-508.
9. Binchun L, Yanqin L, Qiaobin H. Antioxidant activity of flavonoids from tartary buckwheat bran. *Toxicological & Environmental Chemistry. From Toxicological & Environmental Chemistry*. 2015; Ahead of Print.
10. Amit K, Neelesh J, Mehra K, Swarnakar K. Role of Antioxidants for the Treatment of Cardiovascular Diseases. Challenges and Opportunities. *From Current Pharmaceutical Design*. 2015; 21 (30): 4441-4455.
11. KokFookSeng B, Ida I, Lee CT, Firdausi R, Shahrulzaman S. Biologically active quercetin from onion and its effect on health. *Elixir Food Science*. 2015; 78: 30057-30063.
12. Abdel-Gawad Mahfouz M, El-Hashash Maher A, El-Sayed Mortada M, El-Wakil Eman A, Abdel-Lateef Ezzat E. Chromatographic isolation of *Allium cepa* (ssp. red onion) and its cytotoxic activity against human liver carcinoma cell lines (HEPG2). *Int J Pharm Pharm Sci*. 2014; 6 (8): 108-111.
13. Sato A, Zhang T, Yonekura L, Tamura H. Antiallergic activities of eleven onions (*Allium cepa*) were attributed to quercetin 4'-glucoside using QuEChERS method and Pearson's correlation coefficient. *Journal of Functional Foods*. 2015; 14: 581-589.
14. Simin N, Orci D, Cetojevic-Simin D, Mimica-Dukic N, Anackov G, Beara I, Mitic-Culafic D, Bozin B. Phenolic profile, antioxidant, anti-inflammatory and cytotoxic activities of small yellow onion (*Allium flavum* L. subsp. *flavum*, Alliaceae). *Food Science and Technology*. 2013; 54: 139-146.
15. Yang, Anping. An activity ingredient for treating prostatic hyperplasia and its medicine. *From Faming Zhuanli Shenqing*, 2015. CN 105030800 A 20151111.
16. Jiang Q, Yuan Z, Li J, Hou Y. Extraction of β -sitosterol from *Polygonum tinctorium* Aiton and study on its antioxidation. *From Shipin Gongye Keji*. 2015; 36 (6): 108-112.
17. Bagrationi D. Iadigar Daudi. Edition of Tbilisi University, Republic of Georgia, 1993.
18. Mskhiladze L, Chincharadze D, Tits M, Wauters JN, Bellahcène A, Castronovo V, Mouithys-Mickalad A, Franck T, Frédérick M. Identification of Cytotoxic and Antioxidant Compounds from *Allium gramineum* Flowers. *International Journal of Pharmaceutical Sciences and Drug Research* 2015; 7(1): 78-82.
19. Mskhiladze L, Chincharadze D, Mshvildadze V, Pichette A, Frédérick M, Ollivier E, Elias R. Steroidal Glycosides from the Flowers of *Allium leucanthum*. *Chemistry of Natural Compounds*. 2015; 51 (5): 900-9004.
20. Mskhiladze L, Kutchukhidze J, Chincharadze D, Delmas F, Elias R, Favel A. In vitro antifungal and antileishmanial activities of steroidal saponins from *Allium leucanthum* C. Koch a Caucasian endemic species. *Georgian Medical News*. 2008; 1 (154): 39-43.
21. Mskhiladze L, Legault J, Lavoie S, Mshvildadze V, Kuchukhidze J, Elias R, Pichette A. Cytotoxic Steroidal Saponins from the Flowers of *Allium leucanthum*. *J. Molecules*. 2008; 13: 2925-2934.
22. Faizi S, Ali M, Saleem R, Bibi S. Complete ^1H and ^{13}C NMR assignments of stigma-5-en-3-O- β -glucoside and its acetyl derivative. *Magn. Reson. Chem*. 2001; 39: 399-405.
23. Lavoie S, Ouellet M, Fleury PY, Gauthier C, Legault J, Pichette A. Complete ^1H and ^{13}C NMR assignments of a series of pergalloylated tannins. *Magn. Reson. Chem*. 2015; Ahead of Print.
24. Jaramillo K, Dawid C, Hofmann T, Fujimoto Y, Osorio C. Identification of antioxidative flavonols and anthocyanins in *Sicana odorifera* fruit peel. *J. Agric. Food Chem*. 2011; 59: 975-983.
25. Gamal-Eldeen AM, Kawashty SA, Ibrahim LF, Shabana MM, El-Negoumy SI. Evaluation of antioxidant, anti-inflammatory, and antinociceptive properties of aerial parts of *Vicia sativa* and its flavonoids. *J. Nat. Remedies*. 2004; 4: 81-96.
26. Bonaccorsi P, Caristi C, Gargiulli C, Leuzzi U. Flavonol glucosides in *Allium* species: A comparative study by means of HPLC-DAD-ESI-MS-MS. *Food Chem*. 2008; 107: 1668-1673.
27. Soininen TH, Jukarainen N, Julkunen-Tiitto R, Karjalainen R, Vepsäläinen, JJ. The combined use of constrained total-line-shape ^1H NMR and LC-MS/MS for quantitative analysis of bioactive components in yellow onion. *J. Food Compos. Anal*. 2012; 25: 208-214.

Source of Support: Nil, Conflict of Interest: None declared.