



Antibacterial Activity and Phytochemical Study of Ethanolic Extract of *Triumfetta rhomboidea* Jacq

V P Devmurari*, T J Ghodasara, N P Jivani

Smt. R B Patel Mahila Pharmacy College, Bhavnagar Road, Kailashnagar, Atkot, Gujrat, India

ABSTRACT

Ethanolic extract of *Triumfetta rhomboidea* Jacq was subjected to various phytochemical tests. Preparative Thin layer Chromatography study of the extract was performed and active constituents were isolated. Spectral analysis of the isolated constituent indicates that *Triumfetta rhomboidea* (Tiliaceae) contains carbohydrate glycosides, phytosterol, steroids, flavonoids, tannin & phenolic compounds and triterpenoids. Antibacterial activity of ether and alcoholic extract of the plant was performed. Results exhibited that *Triumfetta rhomboidea* Jacq contain good antibacterial action.

Keywords: *Triumfetta rhomboidea* Jacq, Preparative Thin layer Chromatography, Phytochemical.

INTRODUCTION

Triumfetta rhomboidea is a perennial herb having important role in ancient therapy. Various Parts of the plant used therapeutically are fruit, flower, leaves, bark and root. Root is tonic styptic, galactagogue, aphrodisiac, cooling, useful in dysentery and as diuretic. Pounded roots are given in the treatment of Intestinal ulcer. Leaves, Flowers and Fruit are mucilaginous demulcent, astringent, and also used in gonorrhoea and against leprosy.¹⁻⁴ In the present study, active constituents of the plant were analyzed and evaluated for antibacterial activity.

EXPERIMENTAL

Phytochemical Screening

Triumfetta rhomboidea Jacq (Tiliaceae) was procured from botanical garden of B K Mody Govt Pharmacy College, Rajkot. The leaves of *Triumfetta rhomboidea* were dried under shade and powdered with a mechanical grinder. Dried material was extracted with ethanol (90% v/v) in Soxhlet apparatus and after complete extraction (50 hr) the solvent was removed by distillation under reduced pressure and resulting semisolid mass was vacuum dried.⁵⁻¹⁴

Ethanolic extract (EETR) of *Triumfetta rhomboidea* were subjected to preliminary phytochemical screening for the detection of various plants constituent.

Test for alkaloids

***Corresponding author: Devmurari V. P.,**
Smt R.B. Patel Mahila Pharmacy College,
Kailashnagar, Bhavanagar road, Atkot-360040
Ta: Jasadan, Dist: Rajkot India
E-mail: Viraldev1985@gmail.com

The small portion extracts were stored separately with a few drops of dilute hydrochloric acid and filtered. The filtrate was tested with various alkaloidal agents, such as Mayer's reagent (cream precipitate) Dragendorff's reagent (orange brown precipitate)

Test for carbohydrates and glycosides

Small quantity of ethanolic extract was dissolved separately in 5 ml of distilled water and filtered. The filtrate may be subjected to Molisch's test to detect the absence of carbohydrates.

Another small portion of extract was hydrolyzed with dilute hydrochloric acid for few hours in water-bath and was subjected to Liebermann- Burchard's, legal and Borntrager's test to detect absence of different glycosides. (Pink to red color indicates presence of glycosides)

Test for flavonoids

5 ml of dilute ammonia solution were added to a portion of aqueous filtrate of plant extract followed by addition of concentrated H₂SO₄. A yellow coloration absorbed in extract indicated presence of flavonoids.

Test for steroids

2 ml acetic anhydride was added to 0.5 g ethanolic extract with 2 ml H₂SO₄. The color changed from violet to blue or green in samples indicated presence of steroid.

Test for terpenoids (salkowski test)

Five ml of extract was mixed in 2 ml of chloroform, and concentrated H₂SO₄ (3ml), was carefully added to form a layer. A reddish brown coloration of the interface was formed indicated presence of terpenoids.

Test for saponin

About 1 ml of alcoholic and agrees extract was diluted with distilled water to 20 ml and shaken in graduated cylinder for

15 minutes. One cm layer of foam indicated presence of saponin.

Test for tannin

When a drug is treated with vanillin-hydrochloric acid reagent, pink or red color is formed due to formation of phloroglucinol.

Test for protein

Mellon's reaction: Million's reagent (mercuric nitrate in nitric acid containing a trace of nitrous acid) usually yields a white precipitate on addition to a protein solution which turns red on heating.

Test for volatile oil or essential oil

Place a thick section of drug on glass slide. Add a drop of Sudan red 3rd reagent and after two minute wash with 50% alcohol mount in glycerin.

Preparative Thin Layer Chromatography

Ordinarily, microgram quantities of mixture of organic compounds are separated by analytical TLC. It is possible to scale up the quantities to milligram amount (10-50mg) by using thicker layer (0.5-2.0 mm thickness) of the support material and by the use of larger plates (20 × 20 cm or 20 × 40 cm). Multiple developments also bring about better resolution. Preparative TLC for the isolation of marker compound from the ethanolic extract of *Triumfetta rhomboidea* leaves was performed by using solvent system Toluene: Ethyl acetate (9:1).¹⁵⁻²⁰

Antibacterial Activity

In the present research work, the antibacterial activity spectrum of ethanolic extract and ether extract of *Triumfetta rhomboidea* Jacq was analyzed.²¹⁻²⁸ (Table 3 & 4) Three Gram positive bacteria, *Staphylococcus aureus* (MTCC 737), *Enterococcus faecalis* (MTCC 439), *Bacillus cereus* (MTCC 430) and three Gram negative bacteria *Klebsiella pneumoniae* (MTCC 109), *Pseudomonas aeruginosa* (MTCC 2642), *Escherichia coli* (MTCC 1687) were used. Inoculum size was adjusted to 1 to 2 × 10⁷ CFU (Colony Forming Units)/ml by serial dilution with sterilized nutrient broth media. Nutrient agar (pH 7.2-7.4) was used for routine susceptibility testing of nonfastidious bacteria. Stock solution of 10000µg/ml was prepared in 20 % v/v water in DMSO. Using the stock solution, 6000µg/ml, 4000µg/ml, 2000µg/ml and 1500µg/ml solutions were prepared from which 100 µl solution was taken for assay. Ciprofloxacin was used as a standard. 20 % v/v WFI in DMSO was used as a control. Antibacterial assay was carried out by agar Well Diffusion Method. [17-19] After 16 to 18 hours of incubation, each plate is examined.

RESULT AND DISCUSSION

Phytochemical screening suggests that ethanolic extract contain various constituents which are given in the Table 1. Preparative TLC study revealed presence compounds COMP-01, COMP -02, COMP -03, COMP -04 and COMP -5. The compound COMP -01 to COMP -05 gives positive Knollar's and Libermann – Burchred test and the colour produced was typical of triterpenes. IR spectrum produced was similar to triterpenes. IR spectrum in the fundamental region also supported triterpense structure as the bands were noticed due to O-H stretching and C-H stretching of alkanes. (Table 2)

The results of preliminary evaluation showed that *Triumfetta rhomboidea* Jacq possess good antibacterial activity. *P. aeruginosa* and *E. coli* are resistant or less susceptible to *Triumfetta rhomboidea* Jacq.

Table 1: Data showing the preliminary phytochemical screening of the two extracts of *Triumfetta rhomboidea*.

Phytochemical	Presence/Absence
Carbohydrate	++
Glycosides	++
Alkaloids	--
Phytosterol and steroids	++
Flavonoids	++
Protein& Amino Acid.	--
Tannin & phenolic compounds	++
Triterpenoids	++

Table 2: IR and UV Spectral data of the isolated samples.

Code	IR data cm ⁻¹	UV
01	3591 (C-H Stretching in alkenes), 2956 (C-H Stretching in the alkanes.), 1731, 1701, 1683 (C=O Stretching), 1286, 1334 (C-H bending vibration in the alkynes.), 898, 794, 723 (aromatic hydrocarbons.), 1014 (Diphenyl methanol.)	245
02	1201 (O-H stretching in phenol) 3670 (C-H Stretching in alkenes), 2956 (C-H Stretching in the alkanes), 1731,1716,1683 (C=O Stretching), 1222, 1271, 1340 90 (C-H bending vibration in the alkynes), 794, 729, 682 (aromatic hydrocarbons), 1222 (O-H stretching in phenol)	270
03	3151 (C-H Stretching vibration in alkenes.), 2956 (C-H Stretching vibration in the alkanes.), 1745,1735,1683 (C=O Stretching), 1253, 1286 (C-H bending in the alkynes.) 796, 757, 723, 688 (aromatic hydrocarbons) 1253 (O-H stretching)	205
04	3006,3076 (C-H Stretching in aromatic ring), 2956 (C-H Stretching in alkene), 1735,1716,1685 (C=O Stretching), 1224, 1271, 1311 (C-H bending vibration in alkyne), 793, 725 (aromatic hydrocarbons), 1224cm ⁻¹ (O-H stretching {phenolic})	225
05	3672,3735 (C-H Stretching), 2956 (C-H Stretching), 1733,1718,1701(C=O Stretching), 1274, 1311, 1355 (C-H bending vibration), 881, 794, 777, 723, 682 (aromatic hydrocarbons), 1213 (O-H stretching)	295

Table 3. Zone of inhibition of different concentration of ethanolic extract of *Triumfetta rhomboidea* Jacq against test microorganism.

	<i>S. aureus</i>	<i>B. cereus</i>	<i>Ent. faecalis</i>	<i>E. coli</i>	<i>Ps. aeruginosa</i>	<i>Kl. pneumoniae</i>
STD	39.10 ± 0.95	36.67 ± 0.61	30.67 ± 0.61	35.60 ± 0.53	41.07 ± 1.01	36.53 ± 0.61
150 µg/ well	11.13 ± 0.76	11.20 ± 0.20	8.47 ± 0.42	1.00 ± 0.20	0.00	6.20 ± 0.20
200 µg/ well	22.37 ± 0.78	25.20 ± 1.06	21.87 ± 1.20	1.70 ± 0.10	2.47 ± 0.12	21.40 ± 1.25
400 µg/ well	25.33 ± 0.70	27.27 ± 1.10	27.07 ± 0.92	2.40 ± 0.20	2.80 ± 0.20	23.27 ± 1.10
600 µg/ well	28.30 ± 0.95	31.47 ± 1.62	29.73 ± 1.62	3.07 ± 0.12	3.00 ± 0.20	28.87 ± 1.03

Table 4. Zone of inhibition of different concentration of ether extract of *Triumfetta rhomboidea* Jacq against test microorganism.

	<i>S. aureus</i>	<i>B. cereus</i>	<i>Ent. faecalis</i>	<i>E. coli</i>	<i>Ps. aeruginosa</i>	<i>Kl. pneumoniae</i>
STD	39.10 ± 0.95	36.67 ± 0.61	30.67 ± 0.61	35.60 ± 0.53	41.07 ± 1.01	36.53 ± 0.61
150 µg/ well	5.60 ± 0.72	4.33 ± 0.30	8.60 ± 0.53	0.00	0	5.33 ± 0.42
200 µg/ well	9.30 ± 0.75	8.73 ± 0.64	11.67 ± 0.42	0.00	0	8.33 ± 0.31
400 µg/ well	10.13 ± 0.70	10.67 ± 0.61	13.07 ± 0.61	0.00	0	8.47 ± 0.31
600 µg/ well	10.77 ± 0.95	11.67 ± 0.42	14.60 ± 0.60	2.07 ± 0.31	3.20 ± 0.20	10.47 ± 0.42




Fig. 1: Graphical presentation of Inhibition Zone of different concentration of ethanolic extract of *Triumfetta rhomboidea* Jacq against test microorganism




Fig. 2: Graphical presentation of Inhibition Zone of different concentration of ether extract of *Triumfetta rhomboidea* Jacq against test microorganism

REFERENCES

- Chopra R, Nayar S, Chopra I. Glossary of Indian Medicinal Plant, 3rd Edn., Council of Scientific and Industrial Research, New Delhi, 1986, pp 249.
- Mukharjee PK. In, Quality Control of Herbal Drugs, 1st Edn, Business Horizons, Pharmaceutical Publishers, 2002, pp 40.
- Barnes. An Introduction to Herbal Medicinal Products. The Pharmaceutical Journal. 2002; 268: 804.
- Chatterjee A, Chandra Prakash S. The Treatise of Indian Medicinal Plant, Vol 2, National Institute of Science and Communication., CSIR, New Delhi, 1992, pp 170-171.
- Basu BD. Indian Medicinal Plant, Part-I, Dehradun, 1997, pp 159.
- Kirtikar KR, Basu BD. Indian Medicinal Plants, Vol-I, International Book Distributors, India, 2005, pp 76-80.
- Jagetiya GC, Rao SK. Rubia cordifolia - Wound healing property and Free radical scavenging activity. Biol. Pharm. Bull. 2006; 29(3):460.
- Shylesh BS, Nair SA, Subramaniam A. Induction of cell-specific apoptosis and protection. Indian J. Pharmacol. 2005; 37(4): 232.
- Mitra SK, Chatterjee BN, Chakravarthi D, Maiti BL. Chemical investigation of the roots of *Xanthium strumarium*. J Indian Chem Soc. 2006; 83(5): 513-6.
- Osore H. Ecobolic properties of *Triumfetta rhomboidea* Jacq on the gravid mammalian uterus. East Afri. Medi. J. 1982; 59(11): 733.
- Maksoud S, Hosni A, Hanaa HA. The distribution and concentration of urease in the seeds of *Triumfetta rhomboidea* Jacq. Egypt. J. Physio. Sci. 1998; 21(2): 209.
- Nair AGR, Seetharaman TR, Voirin B, Bonvin JF. The true structure of Triumboidin, a flavone glycosides from the leaves of *Triumfetta rhomboidea* Jacq. Phytochemistry 1986; 25(3): 786.
- Kusmi T, Chang CC, Wheeler M, Kubo I, Nakanishi K. Isolation of Triumferol and structural determination by standard spectral method and reported the antigermination activity against lecture seeds from the *Triumfetta rhomboidea* Jacq. Tetra. Lett. 1981; 22(36): 786.
- Pradhan D, Panda PK, Kar DM. Study of antiulcer activity of roots of *Triumfetta rhomboidea*. Journal of Science and Pharmacy 2003; 5: 18-21.
- Chiranjibi PC, Reddy S, Murthy MSR. An ethnobotanical survey of medicinal plants used by the Didayi tribe of Malkangiri district of Orissa, India. Fitoterapia 2008; 79(1): 67-71.
- Mevy JP, Bessiere JM, Dherbomez M, Millogo J, Viano J. Chemical composition and some biological activities of the volatile oils of a chemo type of *Lippia chevalieri*. Food Chemistry 2007; 101(2): 682-685.
- Hansen K., Nyman U, Smit UW, Adersen A, Gudiksen L, Rajasekharan S, Pushpangadan P. In vitro screening of traditional medicines for anti-hypertensive effect based on inhibition of the angiotensin converting enzyme ACE. Journal of Ethnopharmacology 1995; 48(1): 43-51.
- Cohen JH. Fruits and Vegetable Intakes and Prostate Cancer Risk" Journal of National Cancer Institute, 2002; 92: 61-8.
- Pascual C, Gonzalez R, Torricella RG. Scavenging Action of Propolis Extract against Oxygen Radicals. J.Ethnopharmacol. 1994; 41: 9-13.
- Bushman JL. Green Tea and Cancer in Humans; A Review of Literature. Nutr Cancer 1998; 31: 151-159.
- Macho A, Lucena C, Sancho R, Daddario N, Minassi A, Munoz E, Appendino G. Nonpungent Capsaicinoids from Sweet Pepper Synthesis and Evaluation of Chemoprotective and Anticancer Potential. Eur. J. Nutr. 2003; 42(1): 2-9.
- Hasani P, Nargues Sanaz Y, Ghanbari V, Mohammadirad A, Dehghan G, Abdollahi M. In vivo antioxidant potential of Teucrium polium, as compared to α -tocopherol. Acta Pharmaceutica 2007; 57 (1): 24.
- Mevy JP, Bessiere JM, Rabier J, Dherbomez M, Ruzzier M, Millogo J, Viano J. Composition and antimicrobial activities of the essential oil of *Triumfetta rhomboidea* Jacq. Flavour and Fragrance Journal 2005; 21 (1): 80-83.
- Srinivasan KK, Subramaniam SS. Chemical investigation of Emilia sonchifolia. Fitoterapia. 1981; 5: 241-243.
- Manohar K, Adwankar, Manik P. Chitnis. In vivo Anti-Cancer Activity of RC-18 A Plant Isolate from *Rubia cordifolia*, Linn, against a Spectrum of Experimental Tumour Models. Experimental Chemotherapy. 1982; 28: 291-293.
- Gislene GF, Nascimento, Locatelli J, Paulo C. Freitas, Giuliana. Silva. Antibacterial activity of plant extracts and phytochemicals on antibiotic resistant bacteria. Brazilian Journal of Microbiology 2000; 31: 247-256.
- Fisgin NT, Cayci YT, Coban AY, Ozatli D, Tanyel E, Durupinar B, Tulek N. Antimicrobial activity of plant extract ankaferd blood stopper. Fitoterapia 2009; 80: 48-50.
- D. Ayfer atefi, Erdo OT. Antimicrobial activities of various medicinal and commercial plant extracts. Turk J Boil 2003; 27: 157-162.